

Atypical Antidepressants Cheat Sheet

by adrianao via cheatography.com/168237/cs/35240/

Medication

Buproprion

Purpose

Acts by inhibiting norepinephrine and dopamine uptake, and is referred to as a norepinephrine-dopamine reuptake inhibitor.

Therapeutic Uses

Treatment for depression

Alternative to SSRIs and SNRIs for clients unable to tolerate sexual dysfunction adverse effects of these antidepressants

Aid for smoking cessation

Prevention of seasonal pattern depression

Alternative treatment choice for attentiondeficit disorder

Complications

Complications	
Complication	Nursing Action and
	Education
Headache, dry	Treat headache with mild
mouth, GI	analgesic. Observe for
distress,	effects and notify the
Constipation,	provider if intolerable. Sip
Increased heart	on fluids to treat dry
rate, Hypert-	mouth and increase
ension, Restle-	dietary fiber to prevent
ssness, and	constipation.
Insomnia	

Nausea, Vomiting, Anorexia, Weight Loss

Seizures

Monitor weight and food intake

, intak

Avoid administering to clients at risk for seizures (a client who have head injuries). Monitor for seizures, and treat accordingly.

Contraindications/ Precautions

Bupropion is a pregnancy risk category B.

Notify the provider if pregnant or breastfeeding. The provider will determine is the risks of taking the medication out-weigh the benefits.

Contraindicated in clients taking MAOIs.

Contraindicated for clients who have seizure disorders or eating disorders.

Interactions

Interaction	Nursing Action
MAOIs	MAOIs should be discon-
(phenelzine)	tinued 2 weeks prior to
increase the	beginning treatment with
risk for toxicity	bupropion.

Medication

Mirtazapine

Purpose

Referred to as a serotonin-norepinephrine disinhibitor.

It increases the release of serotonin and norepinephrine by blocking the presynaptic receptors, and thereby increases thee amount of neurotransmitters available for impulse transmission.

Nursing Action

Therapeutic effects can occur sooner with less sexual dysfunction than with SSRIs

Mirtazapine is generally well tolerated.
Clients can experience sleepiness that can be exacerbated by other CNS depressants (alcohol, benzodiazepines), weight gain, and elevated cholesterol.

Client Education

Take at bedtime, can be used as sleep aid.

Medication

Vilazodone

Purpose

Both blocks serotonin and works with a serotonin agonist at receptor site (first medication to work in this way).

Nursing Action

Contraindicated with SSRIs and SNRIs (serotonin syndrome), and other serotonin receptor agonists (buspirone and phenothiazines). Stop MAOIs at least 14 days before starting vilazodone.

Teach manifestations of serotonin syndrome to client and instruct them when to notify provider.

Monitor for sucidal ideation

Many adverse effects are similar to those of SSRIs and SNRIs.

Take with food to help increase absorption.

Client Education

Avoid grapefruit juice while taking Vilazodone because grapefruit juice inhibits CYP3A4 metabolism resulting in an increase in the medication blood level.

Medication

Nefazodone

Purpose

Selectively inhibits the reuptake of serotonin and norepinephrine.

Nursing Action

Rapidly absorbed in 1 hr when taken without food.

Adverse effects are sleepiness, headache, dizziness, blurred vision, dry mouth, nausea, constipation, weight gain, and sexual dysfunction

Stop MAOIs at least 14 days before starting nefazodone.

Medication

Trazodone ER

By adrianao cheatography.com/adrianao/ Published 5th November, 2022. Last updated 5th November, 2022. Page 1 of 2. Sponsored by CrosswordCheats.com Learn to solve cryptic crosswords! http://crosswordcheats.com



Atypical Antidepressants Cheat Sheet by adrianao via cheatography.com/168237/cs/35240/

Purpose

Moderate selective blockage of serotonin receptors, which allows more serotonin to be available for impulse transmission.

Nursing Action

Usually used with another antidepressant agent.

Sedation is a potential problem, can be indicated for a client who has insomnia.

Priapism is a potential adverse effect. Instruct clients to seek medical attention immediately if this occurs

Don't drink grapefruit juice as it inhibits CYP3A4 metabolism resulting in an increase in the medication blood level resulting in toxicity.



By adrianao cheatography.com/adrianao/

Published 5th November, 2022. Last updated 5th November, 2022. Page 2 of 2. Sponsored by CrosswordCheats.com
Learn to solve cryptic crosswords!
http://crosswordcheats.com