

Definitions:


DEFINITIONS OF MICROSCOPIC TERMS

Acantholysis	Loss of intercellular cohesion between keratinocytes
Acanthosis	Diffuse epidermal hyperplasia
Dyskeratosis	Abnormal, premature keratinization within cells below the stratum granulosum
Erosion	Discontinuity of the skin showing incomplete loss of the epidermis
Exocytosis	Infiltration of the epidermis by inflammatory cells
Hydropic swelling (ballooning)	Intracellular edema of keratinocytes, often seen in viral infections
Hypergranulosis	Hyperplasia of the stratum granulosum, often due to intense rubbing
Hyperkeratosis	Thickening of the stratum corneum, often associated with a qualitative abnormality of the keratin
Lentiginous	A linear pattern of melanocyte proliferation within the epidermal basal cell layer
Papillomatosis	Surface elevation caused by hyperplasia and enlargement of contiguous dermal papillae
Parakeratosis	Keratinization with retained nuclei in the stratum corneum. On mucous membranes, parakeratosis is normal.

Definitions (cont.):

Spongiosis	Intercellular edema of the epidermis
Ulceration	Discontinuity of the skin showing complete loss of the epidermis revealing dermis or subcutis
Vacuolization	Formation of vacuoles within or adjacent to cells; often refers to basal cell–basement membrane zone area

Skin Diagram:



Major Skin Cells:

- 1) Keratinocytes
- 2) Melanocytes: anchor to basement membrane and pump out pigment for protection from sunlight damage
- 3) Langerhans' cells: antigen-presenting system

Normal Skin Structure and Function:

Function: Barrier between the individual and the external environment

Burn victims:

loss of barrier leading to loss of fluid and secondary bacterial infections.

Eczema/Dermatitis:

damage to the skin resulting in inflammation and spongiosis with edema of the dermis. Hypersensitive reaction to certain chemicals or drugs. Atopic eczema patients are predisposed to allergic rhinitis and asthma.

Immersion:

prolonged immersion in water may overwhelm the barrier (ie. occlusal dressing of burns or under surgical dressings) - can be an advantage in topical therapy.

Incidences of Skin Diseases:

Common: acne, psoriasis, eczema, seborrheic warts and viral warts, actinic raitosis, basal cell carcinoma, squamous cell carcinoma.

Uncommon: pemphigoid, pemphigus, melanoma and scabies.

Rare: xeroderma pigmentosum, mycosis fungoides.

Age, Sex, Anatomical Site,

Rare, Exposure and Geography.

Disorders:

Inflammatory&Hematopoietic cells

Polymorphonuclear

leucocytes (polymorphs)– accumulate in the skin due to response to infections e.g Staphylococcus aureus in impetigo.

Psoriasis: neutrophilic infiltrate with intra-epithelial migration forming sterile abscesses. Others are Sweet's disease and pyoderma gangrenosum show a heavy dermal neutrophilic infiltrate – 2nd to chronic inflammatory bowel disease.

Autoimmune conditions: e.g dermatitis herpetiformis show neutrophilic infiltrate where as bullous pemphigoid shows an eosinophilic infiltrate.

Disorders:

Inflammatory&Hematopoietic cells (cont)

Lymphocytic infiltrate: seen in most chronic inflammatory skin diseases and comprises mostly T-lymphocytes (CD4/Helper T-cells)

Eczema: lymphocytic infiltrate and spongiosis

Lupus erythematosus and lichen planus

Cutaneous lymphomas:

primary (mycosis fungoides) or secondary (peripheral T-cell lymphoma, anaplastic large cell lymphoma etc.)

Mycosis fungoides: T-cell lymphoma – Sézary syndrome

Histiocytic infiltrates: seen in granulomatous skin diseases e.g. bacterial (mycobacteria), fungal (candidiasis) or protozoal infections.

Systemic diseases: e.g. Sarcoidosis.

Primary granulomatous skin diseases: granuloma annulare or necrobiosis lipoidica.

Infections:

Clinical appearance depends on:

1. Site
2. Nature of organism
3. Nature of the body's response to the infection.

Routes of transmission:

hematogenous or penetration of the skin barrier

Viral Infections:

Viruses are obligate intracellular organisms – metabolic basal cells.

Human Papilloma Virus (HPV):

DNA virus with numerous subtypes.

Skin and oral mucosa causes squamous papillomas, verrucae vulgaris, verruca plantaris, verruca plana, condyloma acuminatum (genital warts)

Molluscum contagiosum:

DNA pox virus – umbilicated self-limiting lesion in children.

Herpes virus: HVS1 and HPV2

(blistering vesicles), herpes zoster (chickenpox) and shingles.

Human herpes virus 8 (HHV8):

Kaposi's sarcoma (HIV)

Bacterial Infections:

Impetigo: Staph.aureus in children but Strep. in elderly

Cellulitis: Streptococcus pyogenes

Tuberculosis of the skin (lupus vulgaris): Mycobacterium tuberculosis or Mycobacterium bovis the cause of scrofuloderma.

Bacterial Infections: (cont)

Leprosy (Hansen's disease):

Mycobacterium leprae (lepomatous and tuberculoid). Lepomatous form fatal while tuberculoid form destroy tissue and nerves resulting in mutilated leonine facies and auto-amputations of digits.

Fungal and Protozoa Infections:

Fungal:

Common ringworm (tinea) Tinea pedis (athletes foot) Candida, Blastomyces and Norcardia

Protozoal:

Leishmaniasis infections transmitted by sandflies, organism infiltrates the macrophages

Uticaria:

Hives or wheals (reaction pattern), itching and swelling. Sudden marked increase in the permeability of the dermal blood vessels resulting in edema of the dermis – erythematous or edematous lesions.

Histology: shows marked infiltrate of eosinophils and mast cells

Uticaria: (cont)

Causes: Plant and animal toxins, physical stimuli e.g. heat, cold or stress, various drugs (aspirin and antibiotics)

Histamine is the mediator and increase in IgE.

Lupus erythematosus (SLE or LE)

-Autoimmune disease affecting connective tissue with antibodies directed against DNA.

-Multisystem disease involving almost any organ, most common is skin and kidneys.

-Epidermis and adnexa involved.

-Skin only: Discoid lupus – erythematous, scaly and older lesions show hyperpigmentation.

-Often symmetrical on the face with a butterfly rash over nose and cheeks, and on scalp it may cause scarring alopecia.

-Immunofluorescence show IgG and IgM at the epidermal basement membrane – “lupus band test”

Psoriasis:

Genetically determined – HLA haplotypes (HLA Cw6, B13 and B17).

Disease of epidermal proliferation and excess keratin production, driven by cytokines released from activated T-cells in the dermis Silver-grey scales of parakeratosis on extensor surfaces such as knees and elbows.

First appearance may be at site of trauma e.g. surgical wound – “Koebner effect”.

Result in destructive arthropathy (psoriatic arthritis)

Small bleeding points – Auspitz's sign.

Histology: rete ridges becomes acanthotic with the dermal papillae covered by thin epidermis two or three layers thick.

Loss of the granular cell layer.



By AlbertEinstein

Published 22nd March, 2018.
Last updated 14th March, 2018.
Page 2 of 5.

Sponsored by [Readability-Score.com](https://readability-score.com)
Measure your website readability!
<https://readability-score.com>

Psoriasis: (cont)

Erythematous lesions - caused by dilated vessels in upper dermis. Numerous polymorphs that migrate from the vessels into the epidermis forming pustules (micro-abscesses)- pustular psoriasis. May also involve the sole of feet and palms of hands.

Treatment: Coal tar, Methotrexate and arsenic.

Current therapy: Retinoid (Vit A analogues).

Panniculitis:

Inflammation of the subcutaneous fat.

Most common type is **Erythema nodosum** which occurs as painful red nodules on the shins.

Follow a streptococcal infection or in association with inflammatory bowel disease or tuberculosis.

Lichen Planus:

- Characterized by destruction of keratinocytes, probably mediated by interferon-gamma and tumour necrosis factor from T-cell in the dermis.

- Affects the skin, most commonly the inner surface of the wrist.

- Appears as white lacy lesion – presents as itchy, polygonal, violaceous papules that may form blisters.

- May be caused by certain drugs or medications.

- Mucosal surface white striae – Wickham's striae.

- Histology shows a lymphohistiocytic infiltrate in a band-like fashion at the dermo-epidermal junction- classical lichenoid reaction pattern.

- Basal cell vacuolar degeneration, apoptosis, keratinocyte necrosis.

- In contrast to psoriasis, LP show an increase in the granular cell layer.

- Rete ridges show "sawtooth" morphology.

- Treatment: Steroids

Benign epidermal neoplasms & tumour-like:

Skin tags or fibro-epithelial

polyps: seen frequently in the elderly and common in the axilla. Due to friction rather than a true neoplasm.

Seborrheic keratosis (basal cell papillomas):

common in the elderly, dark greasy looking nodules with an irregular surface. Rarely turn malignant. Stuck-on appearance.

Histology: Convulated surface with keratin tunnels (horn cysts). They may become inflamed. Often removed for cosmetic purposes and to exclude melanoma.

Squamous papilloma: Benign neoplasm of squamous epithelium, HPV induced lesion.

Cyst: Epidermal (infundibular) cyst and pilar (tricholemmal) cyst are the most common.

Pre-malignant lesions:

Actinic keratosis: In sun-exposed areas, show dermal solar elastosis, need to evaluate for keratinocyte intra-epithelial neoplasia (KIN I-III)

Bowen's disease: Squamous cell carcinoma in-situ.

Basal cell carcinoma (BCC):

Very common skin malignancy, most commonly on the face of elderly people.

Related to chronic sun exposure.

Locally very aggressive and destructive, however metastasis are extremely rare.

Nodular or superficial BCC has a better prognosis while micro-nodular and morpheic are more aggressive and recurrence is high.

Associated with mutations in the Drosophila gene patched- PTCH1 (tumour suppressor gene) which is a member of the sonic hedge-hog pathway.

Clinically: ulcerated irregular lesions – rodent ulcers with raised pearly borders and blood vessels visible on the border.

Histology: cells look like normal basal epithelial cells, islands of basaloid cells with peripheral palisading.

May develop from the basal cell layer or hair follicles.



By AlbertEinstein

Published 22nd March, 2018.

Last updated 14th March, 2018.


Page 3 of 5.

Sponsored by **Readability-Score.com**

Measure your website readability!

<https://readability-score.com>

Squamous cell carcinoma (SCC):



- Are common and usually caused by chronic UV exposure.
- Other: immunosuppression, irradiation, chemical carcinogens, HPV infections and chronic infections.
- More common in the elderly .
- Rarely SCC arises at the edge of a chronic skin ulcer (Majolin's ulcer).
- Very aggressive and invasive however metastasize late.
- Excision can be curative and SCC are more sensitive to radiation than BCC's.

Keratoacanthoma:

Rapidly growing, cup-shaped squamous epidermal lesion

Involves sun-exposed skin of elderly - face

Keratoacanthoma: (cont)

Regresses spontaneously if left untreated

Histology:

- Endophytic-endophytic squamous proliferation,
- Cup-shaped lesion
- Crater-like center filled with laminated keratotic material
- Resembles a well differentiated squamous cell carcinoma

Melanocyte Derived Lesions:

- > Melanocytes are pigmented cells
- > Found in basal layer of epidermis
- > Melanin is synthesized by melanosomes
- > Found on skin and mucosal surfaces – oral cavity, vagina, conjunctiva etc.

Types of Benign Lesions:

Lentigos:

Increase in single melanocytes in basal areas. Small, pigmented macule.

Naevi:

Increase in groups of melanocytes. Melanocytes form nests.

Types of Benign Lesions: (cont)

Freckle:

Increase in melanin production by normal melanocytes which are taken up by adjacent keratinocytes. UV stimulation.

Naevus:

Common in light-skinned individuals. Absent at birth and appear in early childhood. Increase in number through early adulthood. Found mainly on sun-exposed areas. Different stages

STAGE 1 - JUNCTIONAL:

At dermo-epidermal junction –intraepidermal nests. Tan or brown pin-point macules (1-2mm). Gradually enlarge. Increase in number of individual melanocytes at dermo-epidermal junction.

STAGE 2 – COMPOUND:

Nevus is elevated above skin surface. Childhood \ adolescence. Slightly raised due to nests of melanocytes

Naevus: (cont)

STAGE 3- INTRADERMAL:

Middle to old age. Nevus cells detach from dermo-epidermal junction. Smaller, more mature and less metabolically active. Cannot divide. Two components Junctional and intradermal. Pink due to lack of melanin

Histology of Stage 2:

- Nests of melanocytes at dermo-epidermal junction and infiltrating dermis
- Pigmented
- Dendrites lost
- Nuclei become round, inconspicuous nucleoli

Blue Naevus:

- Occurs in deep dermis
- Bluish tinge
- Any area of skin
- Less than 5mm in diameter
- Solitary
- Malignant transformation is rare

MALIGNANT LESIONS:

Melanoma

- Common skin cancer
- Can arise from any melanocyte, composed of malignant melanocytes.
- Usually pigmented but may be unpigmented.



By AlbertEinstein

Published 22nd March, 2018.
Last updated 14th March, 2018.
Page 4 of 5.

Sponsored by [Readability-Score.com](https://readability-score.com)
Measure your website readability!
<https://readability-score.com>

MALIGNANT LESIONS:

Melanoma (cont)

- Etiology associated with fair skin and sunburn
- Asymmetrical (ABCDE)
- Irregular borders
- Varying colours: blue, black-brown, ulcerated
- >6mm
- Evolution
- Prognosis depends on thickness of lesion and presence of surface ulceration – Breslow thickness
- > 1 mm depth indicates significant risk for metastasis
- Cure rate for completely excised non-ulcerated melanomas below 1mm is 100%

Main Variants:

- 1. Lentigo maligna melanoma:** - sun damaged skin of the elderly, develops from a pre-existing in situ lesion termed a lentigo maligna (Hutchinson's melanotic freckle).
- 2. Acral lentiginous melanoma:** - palms and soles, most common type in Non-Caucasians.
- 3. Superficial spreading melanoma:** - most common type in people from European descent
- 4. Nodular melanoma:** - retain no features to identify a pre-existing in situ lesion

Vesiculo-Bullous Conditions:

Fluid –filled cavities

Within the skin

Caused by the separation of two layers of tissue and leakage of plasma into the space

Bullae: > 5mm

Vesicles: < 5mm

DISTINCT MECHANISMS OF BLISTER FORMATION:

Direct destruction of bonds between epithelium e.g. pemphigus

Cells forced apart by edematous fluid e.g. eczema

Cellular destruction leaving gaps e.g. herpes infection

Basement membrane or its attachments to epidermis altered e.g. bullous pemphigoid

Pemphigus:

BONDS BETWEEN EPITHELIUM IS BROKEN

Common in middle-aged to elderly

Mortality rate of approximately 40%

Autoantibodies directed against desmosomes

Pemphigus: (cont)

Bridges lysed and epidermis falls apart

- Blister formed containing epithelial cells within cavity (acantholysis)

- Various forms:
Pemphigus vulgaris
Pemphigus foliaceus
Pemphigus vegetans

- Skin very fragile
- Firm pressure on normal-looking skin will cause blister formation (Nikolsky's sign)

Bullous Pemphigoid (BP):

More common than pemphigus

Usually over 60 years of age

Self-limiting

Associated with periods of pruritis

Blister forms at dermo-epidermal junction

Etiology:

- > Circulating antibodies against lamina lucida of the BM.
- > Linear deposition of antibodies along basement membrane (IgG)
- > Causes antigen-antibody complexes and release of complement factors as well as degranulation of mast cells
- > Blisters are more persistent

DERMATITIS HERPETIFORME:

- Characterized by small, itchy blisters
- On extensor surfaces of knees and elbows
- Young adults
- May be associated with Coeliac disease
- Pruritic lesions
- Bullous forms at dermo-epidermal junction
- **Immunofluorescence:** Granular deposits of IgA
- Therapy: Response to dapsone

Bullous and Acantholytic Dermatitis:

Intra-epidermal/Suprabasal Cleft:

1. Pemphigus Vulgaris: IgG (intercellular)
2. Pemphigus Foliaceus: IgG (subcorneal)
3. Pemphigus Erythematous: IgG

Sub-epidermal Cleft:

1. Bullous Pemphigoid: linear IgG and C3 at basement membrane.
2. Dermatitis Herpetiformis: granular IgA papillary dermis.
3. bullous Systemic Lupus Erythematous: IgG, IgA and C3.



By AlbertEinstein

Published 22nd March, 2018.

Last updated 14th March, 2018.

Page 5 of 5.

Sponsored by [Readability-Score.com](https://readability-score.com)

Measure your website readability!

<https://readability-score.com>