Formula

speed	s = d/t -> 50 + 30 = 80 miles, 1+1 = 2h -> 80 miles/2h = 40 mph
velocity	v = ∆x/t
average velocity	v avg = ∆x/∆t = xf − xi/tf − ti -> 100 m in 10.61 s -> xf = 100 m, xi = 0 m, tf = 10.61 s, ti = 0 s -> v avg = 100 m - 0 m / 10.61 s - 0 s = 100/10.61 = 9.43 m/s

Speed and Velocity

speed	the distance traveled per unit of time. Speed is a scalar, a quantity that is described by magnitude alone. Constant speed refers to a fixed distance per unit of time. Average speed includes the total distance and total time.

Speed and Velocity (cont)

velocity	the displacement of an object per unit of time. Since displacement includes a direction, so does velocity. Speed with direction. Velocity is a vector a quantity that has both magnitude and direction.
vector	a quantity that has both magnitude and direction
reference frame	the position from which an event is observed
motion map	an image that represents the position, velocity, and acceleration of an object at one-second intervals
scalar	a quantity that is described by magnitude alone
Motion and reference frame	All motion is relative. It depends on a reference frame. An object may appear to move faster or slower depending on the reference frame.

Speed and Velocity (cont)

average velocity	The slope of a line changes when the velocity of an object changes -> The steeper the slope, the greater the velocity. The average velocity will be different than any of the other. Any point on the line will give only an instantaneous velocity.
change in direction	A change in direction is represented when the line on a position-time graph changes from a positive slope to a negative. slope or from a negative slope to a positive slope. A negative slope indicates an object moving towards the origin. A positive slope indicates an object moving away from the origin.
No motion	horizontal line - means object is not moving -> The object's position does not change
Motion	Displayed in a vector !

By **anjuscha**
cheatography.com/anjuscha/

Published 1st October, 2020.
Last updated 3rd October, 2020.

Sponsored by **CrosswordCheats.com**
Learn to solve cryptic crosswords!
http://crosswordcheats.com

Key concepts

Review: Key concepts

	Horizontally Launched Projectile	Projectile Launched at an Angle
Horizontal Motion	$v_x =$ constant $\Delta x = v_x \Delta t$	$v_x = v_{ix} = v_i \cos\theta =$ constant $\Delta x = (v_i \cos\theta)\Delta t$
Vertical Motion	$v_y = a_y \Delta t$ $v_y^2 = 2a_y \Delta y$ $\Delta y =$ $\frac{1}{2}a_y$ $(\Delta t)^2$	$v_{fy} = v_i \sin\theta + a_y \Delta t$ $v_{fy}^2 = v_i^2($ $\sin\theta$ $)^2 + 2a_y \Delta y$ $\Delta y = (v_i \sin\theta)\Delta t + \frac{1}{2}a_y (\Delta t)^2$

Horizontal motion example

EXAMPLE

Riley and Miguel are playing catch. Riley throws the ball at an angle of 25° relative to the ground at a speed of 23.0 m/s. The ball travels 42.0 m to Miguel, who catches the ball. How long was the ball in the air?

Given:

$\theta =$ 25°

$v_i = 23.0$ m/s

$\Delta x = 42.0$ m

Unknown:

$\Delta t = ?$

The equation that we need to use is:

$\Delta x = ($ $v_i \cos\theta$ $) \Delta t$

$\Delta t = \frac{\Delta x}{(v_i \cos\theta)}$

$= \frac{42.0 \text{ m}}{(23 \text{ m/s})(\cos 25°)}$

$=$ 2.0 s

Formula

Two vectors added at a right angle (90°)	$R^2 = A^2 + B^2$
Magnitude and sign of component vectors	$A_x = A\cos\theta$ $A_y = A\sin\theta$
Magnitude of the resultant vector	$R^2 = R_x^2 + R_y^2$
Angle or direction of the resultant vector	$\tan\theta =$ $\frac{R_y}{R_x}$

Displacement during constant acceleration

Displacement during constant velocity	$\Delta x = vt$
Displacement during acceleration	$\Delta x = \frac{1}{2}(v_f - v_i)t$
Total displacement is the sum of the two	$\Delta x = v_i t + \frac{1}{2}($ v_f $- v_i)t$
Terms are combined	$\Delta x = \frac{1}{2}(v_f + v_i)t$
When the initial position is not zero	$x_f = x_i + \frac{1}{2}(v_f + v_i)t$

Acceleration example

Example: A race car moving west speeds up from 17 m/s to 47 m/s in 2 seconds. What is the car's acceleration?

$a = \frac{47 \text{ m/s} - 17 \text{ m/s}}{2 \text{ s}} =$ $\frac{30 \text{ m/s}}{2 \text{ s}}$

$a = 15 \text{ m/s}^2$

$v_i = 17$ m/s

$v_f =$ 47 m/s

$t = 2$ s

acceleration

$a = \frac{\Delta v}{\Delta t} = \frac{v_f - v_i}{t_f - t_i}$


Acceleration

positive acceleration	an increase in velocity over time
negative acceleration	a decrease in velocity over time
acceleration	the rate at which velocity changes over time
constant	staying the same; unchanging
Positive acceleration	speeds up in the positive direction. slows down in the negative direction
Negative acceleration	slow down down in the positive direction. speeds up in the negative direction.
Slope	of the line on a velocity vs. time graph represents acceleration. Positive slope = acceleration, negative slope = negative acceleration

vectors


quadrant	a quarter of the coordinate plane
components	the two parts of a vector that are perpendicular to each other
resultant vector	the sum of two or more vectors
vector resolution	the process by which the components of a vector are determined
Properties of a vector	A vector is a quantity that has both magnitude and direction. Examples of vectors: Displacement, velocity, acceleration. Vectors are drawn using an arrow

More



Sponsored by CrosswordCheats.com
Learn to solve cryptic crosswords!
http://crosswordcheats.com

Sign of a component



The sign of a component depends on the quadrant of the coordinate system it is in.

Projectile Motion

projectile	an object that is set in motion following a path in which the only force acting on it is gravity.
inertia	the natural tendency of objects to resist a change in motion
projectile motion	the curved motion that results from the combination of an object's horizontal inertia and the force due to gravity pulling the object downward. I.e. A ball rolling of the table, A player shooting a jump shot -> Projectiles follow a parabolic path
parabolic	having the shape of a parabola

Projectile Motion (cont)

vectors	Vectors are used to describe motion in two dimensions. Vectors can be broken down into x and y components. The components of a vector are the two parts of a vector that are perpendicular to each other

Add


$\cos\theta = \frac{A_x}{A}$

$\sin\theta = \frac{A_y}{A}$


If we rearrange these we now get:

$A_x = A\cos\theta$


$A_y = A\sin\theta$




Horizontal



Horizontal example



continued



Sponsored by **CrosswordCheats.com**
Learn to solve cryptic crosswords!
http://crosswordcheats.com