

Case

- 80 y.o., pensioner
- Pain in R leg
- Onset 4 months ago
- Pain comes on about half way through his walk to shop (400m), eases off if he sits for a while, then able to resume
- Same thing happens on way back
- 'Cramp-like' sensation
- **ABPI** (diagnosis PAD): 0.8 (results last week)

AF: walking

RF: sitting

AA: walking to shop

Extras

- Hypertension since 55 y.o.
- Hypercholesterolemia diagnosed 10 yrs ago
- Coronary angioplasty 2 yrs ago (following episodes of angina)
- Aspirin (300mgs/day; NSAID)
- Atorvastatin (prevents cardiovascular disease in high risk pts)
- Lisinopril (treats high BP)
- Smoked 20/day since teens
- Drinks pint of ale most nights
- Lives alone (wife died 2 yrs ago)
- 2 children who he doesn't see often
- Doesn't enjoy eating much anymore now he is on his own

Physical Examination Findings

BP: R+L 160/100

General observations

- Looks underweight
- Feet look pale & feel cool to the touch
- No swelling in legs

Skin

- Nicotine staining on fingers from smoking

Chest, cardiovascular & respiratory

- Bruit (partially occluded artery) on the R femoral artery

Peripheral inspection & pulses

- Diminished dorsals pedis * posterior tibia's pulses in the R leg

Clinical tests

- Good ROM at the hips & knees
- Full Lx ROM
- Feels muscle stretch behind knees on full flexion

Discussion

Working diagnosis

- Peripheral Arterial Disease (PAD)
- Non-MSK
- Vascular w/ key Hx findings being **cramping** pain
- **AF**: set distance
- **RF**: sitting
- ABPI: **0.8**, which supports working Dx
- Several risk factors for PAD including **existing CV disease** & being a **heavy smoker & drinker**
- Feet look **pale** & are **cool**: suggests that PAD is more advanced than the ABPI suggests
- **Diminished pulses** in LL w/ **bruit** over R femoral artery

Yellow flags

- High blood pressure
- Underweight
- Nutritional status
- Limited social interactions / social isolation
- Possible depression

Learning Outcomes

Understand Peripheral Arterial Disease (PAD)

- Describes **stenosis/occlusion** of **peripheral arteries**, affecting blood supply to extremities, usually LL
- Commonly caused by **atherosclerosis**
- Acute limb ischemia:**
 - Sx & Ssx develop over <2 weeks
 - **Sudden** onset leg pain
 - Sudden deterioration of **claudication**, associated **loss of pulses** &/or **pallor**
 - **Coldness/cyanosis** of limb
 - **Loss of motor/sensory** functions
- Chronic limb ischemia:**
 - Progressive development of **cramp-like** pain in calf, thigh or buttock (atypical terms: tired, giving way, sore, hurts - rather than describing cramp)
 - LL pain on **walking/exercise**, relieved by **rest**
 - Unexplained leg/foot pain
 - **Non-healing** wounds to LL
 - ABPI 0.0-0.4
- Risk factors:**
 - Smoking
 - Drinking
 - Hx of CV disease
 - Diabetes mellitus



By bee.f (bee.f)
cheatography.com/bee-f/

Published 23rd February, 2024.
Last updated 15th May, 2023.
Page 1 of 4.

Sponsored by CrosswordCheats.com
Learn to solve cryptic crosswords!
<http://crosswordcheats.com>

Learning Outcomes (cont)

Differentiate vascular from neurogenic claudication

Vascular claudication:

- Caused by **reduced** blood flow to LL
- **Sx & Ssx:**
- pain in calves, thighs, or buttocks
- Pain during walking/exercise
- Relieved by rest
- 'Cramping' or 'burning' sensation
- Legs may feel cold, numb, or weak
- Visible signs of poor circulation: skin may appear paler blue, slow wound healing
- Absent/poor LL pulses

Neurogenic claudication:

- Caused by nerve compression/damage in lower back
- **Sx & Ssx:**
- Pain in lower back, buttock, legs (above knees usually)
- Aggravated by standing & walking
- Relieved by sitting or leaning forward
- 'Deep ache' or 'numbness' sensation
- Associated tingling, weakness, shooting pains in legs

Investigations:

- **Ankle-brachial index test (ABPI):** evaluate blood flow to the legs
- **Electromyography (EMG) or nerve conduction studies:** evaluate nerve function
- **MRI or CT:** evaluate the spine & surrounding structures

Differentiate PAD from other vascular disorders

PAD:

- Pain in the calves, thighs, buttocks
- Pain during walking/exercise
- Improves w/ rest
- Other Sx: numbness, weakness, coldness
- Skin may appear pale, shiny, discoloured, cool to touch
- Hair loss/slow growth
- Slow wound healing
- Ulcers/sores

DVT:

- Pain unilaterally
- Other Sx: swelling, redness, warm area
- Aggravated by standing/walking
- May not improve w/ rest
- Skin may appear discoloured, veins visible on skin surface
- Risk: Hx immobility, surgery, family Hx of clots

Chronic Venous Insufficiency (CVI):

- Pain, fatigue, heaviness in legs
- Swelling, possible varicose veins
- Skin appeared thickened, discoloured
- Pain worse w/ prolonged standing/sitting, may improve w/ leg elevation
- Risk: pregnancy, family Hx of varicose veins

How to interpret ABPI?

- <0.5: suggests severe arterial disease
- 0.6 - 0.7: suggest presence of arterial disease or mixed arterial/venous disease
- 0.8 - 1.3: suggest no evidence of significant arterial disease
- >1.3: may suggest presence of arterial calcification (such as pts w/ diabetes, RA, systemic vasculitis, atherosclerotic disease, & advanced chronic renal failure)

Differentiate typical from atypical cramp

Typical cramp:

- Often occur in legs (esp. calves), feet, hands
- AF: during/after exercise & may be related to muscle fatigue or dehydration
- RF: stretching or massage
- Last up to 10 min
- Doesn't usually occur frequently or interfere significantly w/ ADL

Atypical cramp:

- In any muscle, including those not typically affected by cramps
- Occur without obvious trigger / after minimal activity
- Accompanied by other Sx: weakness, stiffness, twitching
- May last longer, & feel more severe
- May occur frequently or interfere w/ ADL

AECC clinic risk management for hypertension

- <140/90: no action required
- 140/90 - <160/100: BP to be measured at next follow-up appt; Letter to GP within 1 week if still >140/90
- 160/100 - <180/110: Tell pt to see GP within 1 week; Tutor send letter to GP within 48hrs; BP to be measured at 1st follow-up appt
- >180 systolic OR >110 diastolic: Floor tutor check BP manually; tutor tell pt to see GP the sea day; Tutor to follow up GP letter same day

Screening questions fro depression

1. During the last **month** have you often been feeling down, depressed or hopeless?
 2. During the last **month** have you often been **bothered** by having little interest or pleasure in doing things?
- IF ANSWER 'YES' TO ONE/BOTH, FOLLOW UP WITH:
- *During the last month, have you often been bothered by:*
3. Feeling **bad** about yourself or that you're a failure or have let yourself or your family down?
 4. Poor **concentration**?
 5. **Tiredness**/ low energy levels?
 6. Changes in **appetite** (reduced or increased)?
 7. Changes in your **sleep pattern** (sleeping too much, problems getting to sleep, waking in the night or waking early)?
 8. Being so **slowed down**, or so **restless/fidgety**, that other people have noticed?
 9. Thoughts of **death**?

