

| Series | | | | |
|--------------------|--|--|---|---|
| Series Type | General Summation | Convergence | Divergence | Notes |
| Infinite Series | $\sum_{n=k}^{\infty} a_n$, where k is some whole number ($k = \{0, 1, 2, \dots\}$) for which a_n is well-defined | Converges if $\lim_{n \rightarrow \infty} S_n = L$, where L is finite | Diverges if $\lim_{n \rightarrow \infty} S_n = \infty$ or DNE | S_n is the partial sum of the series: $S_n = a_1 + a_2 + a_3 + a_4 + \dots + a_{n-1} + a_n$ |
| Harmonic Series | $\sum 1/n$ | Never converges | Always diverges | The alternating version of this series ($\sum (-1)^{n+1}/n$) converges, and $\sum 1/n$ is a P-Series with $p=1$ |
| Geometric Series | $\sum_{n=0}^{\infty} ar^n = \sum_{n=1}^{\infty} ar^{n-1}$, where a is the first term of the series, and r is the common ratio between terms | Converges if $ r < 1$ | Diverges if $ r \geq 1$ | If the series converges, its sum is $S = a/(1-r)$ |
| P-Series | $\sum 1/n^p$, where p is a positive number | Converges if $p > 1$ | Diverges if $p \leq 1$ | |
| Alternating Series | $\sum (-1)^n b_n$, $\sum (-1)^{n+1} b_n$, or $\sum (-1)^{n-1} b_n$ | Converges if $\lim_{n \rightarrow \infty} b_n = 0$ and b_n is a decreasing sequence ($b_{n+1} \leq b_n$ for all n) | Cannot show divergence, inconclusive | |
| Telescoping Series | $\sum (b_n - b_{n+1})$ | Converges if $\lim_{n \rightarrow \infty} S_n = L$, where L is finite | Diverges if $\lim_{n \rightarrow \infty} S_n = \infty$ or DNE | S_n is the partial sum of the series: $S_n = \sum_{i=1}^n (b_i - b_{i+1})$ where n is finite |

Alternating Series Estimation Theorem: If $S_n = \sum_{i=1}^n (-1)^i b_i$ or $\sum_{i=1}^n (-1)^{i-1} b_i$ is the sum of an alternating series that converges, then $|R_n| = |S - S_n| \leq b_{n+1}$

Trigonometric functions like $\cos(n\pi)$ or $\sin(n\pi + \pi/2)$ act as sign alternators, like $(-1)^n$

The Alternating Series Test (AST) checks the limit, but since the AST only concludes convergence, we explicitly apply the Test For Divergence (albeit, redundantly) if the limit fails (even though it checks the same limit)

| Series Tests | | | | |
|---------------------|----------------------------|---------------------------------------|---|-------|
| Test Type | Typical series to use test | Convergence | Divergence | Notes |
| Test for Divergence | $\sum a_n$ | Cannot show convergence, inconclusive | Diverges if $\lim_{n \rightarrow \infty} a_n \neq 0$ or DNE | |



Series Tests (cont)

| | | | |
|---|---|---|---|
| Integral Test | $a_n = f(n)$, which is a positive, continuous, decreasing function on the interval $[k, \infty)$, usually with clearly-integrable functions | Converges if $\int_k^\infty f(n)dn$ converges | Diverges if $\int_k^\infty f(n)dn$ diverges |
| (Direct) Comparison Test | a_n and b_n are positive-termed ($a_n \geq 0$ and $b_n \geq 0$ for all n) and $a_n \leq b_n$ for all n | $\sum a_n$ converges if $\sum b_n$ converges | $\sum b_n$ diverges if $\sum a_n$ diverges Inconclusive if b_n diverges in proving convergence, or a_n converges in proving divergence |
| Limit Comparison Test | a_n and b_n are positive-termed, and $\lim_{n \rightarrow \infty} a_n/b_n = c$, where $0 < c < \infty$ | $\sum a_n$ converges $\iff \sum b_n$ converges | $\sum a_n$ diverges $\iff \sum b_n$ diverges Inconclusive if $c=0$, $c=\infty$, or c DNE |
| Ratio Test | $\sum a_n$, usually with $n!$ terms, product terms, or $(a_n)^n$ | Absolutely converges if $\lim_{n \rightarrow \infty} a_{n+1}/a_n < 1$ | Diverges if $\lim_{n \rightarrow \infty} a_{n+1}/a_n > 1$ Inconclusive if $\lim_{n \rightarrow \infty} a_{n+1}/a_n = 1$ |
| Root Test | $\sum a_n$, usually with $(a_n)^n$ | Absolutely converges if $\lim_{n \rightarrow \infty} a_n ^{1/n} < 1$ | Diverges if $\lim_{n \rightarrow \infty} a_n ^{1/n} > 1$ Inconclusive if $\lim_{n \rightarrow \infty} a_n ^{1/n} = 1$ |
| Absolute/Conditional Convergence Classification | $\sum a_n$ | Absolutely converges if $\sum a_n $ converges | Diverges if $\sum a_n$ diverges Conditionally converges if $\sum a_n $ diverges, but $\sum a_n$ converges |

For the series listed, assume each series to be an infinite series starting at $n=k$: $\sum_{n=k}^\infty = \sum$ as previously defined

If a test is inconclusive, use another test

The symbol $[\iff]$ represents the relationship "if and only if" (often abbreviated to "iff"), meaning both sides of the statement must be true at the same time, or false at the same time

Special Series

| Series | Summation Form | First five terms | Radius & Interval of Convergence |
|--|---|---|--|
| Power Series centered at a | $\sum C_n(x-a)^n$ | $C_0 + C_1(x-a) + C_2(x-a)^2 + C_3(x-a)^3 + C_4(x-a)^4 + \dots$ | $(a-R, a+R)$, $[a-R, a+R)$, $(a-R, a+R]$, or $[a-R, a+R]$ |
| Taylor Series centered at a | $\sum \frac{f^{(n)}(a)(x-a)^n}{n!}$ | $f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \frac{f'''(a)(x-a)^3}{3!} + \frac{f^{(4)}(a)(x-a)^4}{4!} + \dots$ | $ x-a < R$ |
| Maclaurin Series (Taylor Series centered at 0) | $\sum \frac{f^{(n)}(0)(x-0)^n}{n!} = \sum \frac{f^{(n)}(0)x^n}{n!}$ | $f(0) + f'(0)x + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \frac{f^{(4)}(0)x^4}{4!} + \dots$ | $ x < R$ |
| $1/(1-x)$ | $\sum x^n$ | $1 + x + x^2 + x^3 + x^4 + \dots$ | $R=1$, $I=(-1, 1)$ |



Special Series (cont)

| | | | |
|---------------------|--|--|---------------------------------|
| e^x | $\sum x^n/n!$ | $1+x+x^2/2!+x^3/3!+x^4/4!+\dots$ | $R=\infty, I=(-\infty, \infty)$ |
| $\ln(1+x)$ | $\sum (-1)^n x^{n+1}/(n+1)$ | $x-x^2/2+x^3/3-x^4/4+x^5/5-\dots$ | $R=1, I=(-1, 1]$ |
| $\arctan(x)$ | $\sum (-1)^n x^{2n+1}/(2n+1)$ | $x-x^3/3+x^5/5-x^7/7+x^9/9-\dots$ | $R=1, I=[-1, 1]$ |
| $\sin(x)$ | $\sum (-1)^n x^{2n+1}/(2n+1)!$ | $x-x^3/3!+x^5/5!-x^7/7!+x^9/9!-\dots$ | $R=\infty, I=(-\infty, \infty)$ |
| $\cos(x)$ | $\sum (-1)^n x^{2n}/(2n)!$ | $1-x^2/2!+x^4/4!-x^6/6!+x^8/8!-\dots$ | $R=\infty, I=(-\infty, \infty)$ |
| $(1+x)^k$ | $\sum \binom{k}{n} x^n = \sum ((k(k-1)(k-2)\dots(k-n+1))/n!) x^n$ | $1+kx+k(k-1)x^2/2!+k(k-1)(k-2)x^3/3!+k(k-1)(k-2)(k-3)x^4/4!+\dots$ | $R=1$ |
| Taylor's Inequality | $ R_n(x) \leq M x-a ^{n+1}/(n+1)!$, where M is a constant such that $ f^{(n+1)}(x) \leq M$ for all $ x-a \leq d$ | | |

For the series listed, assume each series to be an infinite series starting at $n=0$: $\sum_{n=0}^{\infty} = \Sigma$ as previously defined

The Radius of Convergence, R, is typically found by using the Ratio Test or Root Test

$\binom{k}{n}$ is the "binomial coefficient" (read as "k choose n"). $\binom{k}{n} = k!/(n!(k-n)!)$

$f^{(n)}$ means "the nth derivative of the function f"

$n! = n(n-1)(n-2)\dots(1)$

$n! = n(n-1)(n-2)(n-3)\dots \cdot 3 \cdot 2 \cdot 1$

$0! = 1, 1! = 1$

Areas of Functions

Between two functions $\int_a^b ((\text{top function}) - (\text{bottom function})) dx$

Enclosed by a polar function $\frac{1}{2} \int_a^b f(\theta)^2 d\theta$

Between two polar functions $\frac{1}{2} \int_a^b ((\text{outer polar function})^2 - (\text{inner polar function})^2) d\theta$

Area enclosed by a polar function is with respect to the pole, which is the origin

Average value of a function: $f_{avg} = 1/(b-a) \int_a^b f(x) dx$

Volumes of Solids of Revolution

Disk $\pi \int_a^b (\text{radius})^2 dx$

Washer $\pi \int_a^b ((\text{outer radius})^2 - (\text{inner radius})^2) dx$

Cylindrical Shell $2\pi \int_a^b (\text{radius})(\text{height}) dx$

For Cylindrical Shells: radius=x or y, and height=f(x) or g(y)

Arc Lengths

Surface Areas

Function revolved about an axis $2\pi \int_a^b (\text{radius})(\text{Arc Length component}) ds$

Function revolved about y-axis $2\pi \int_a^b x \sqrt{(f'(x))^2 + 1} dx$

Function revolved about x-axis $2\pi \int_a^b y \sqrt{(g'(y))^2 + 1} dy$

Parametric function of t revolved about y-axis $2\pi \int_a^b f(x) \sqrt{(x'(t))^2 + (y'(t))^2} dt$

Parametric function of t revolved about x-axis $2\pi \int_a^b g(y) \sqrt{(x'(t))^2 + (y'(t))^2} dt$

$f(x)=dy/dx, g(y)=dx/dy, x'(t)=-dx/dt, \text{ and } y'(t)=dy/dt$

Integration by Parts

Indefinite Integral $\int u dv = uv - \int v du$

Definite Integral $\int_a^b u dv = uv|_a^b - \int_a^b v du$

Integration by Parts is used to integrate integrals that have components multiplied together in their simplest form, often referred to as a "product rule for

Trigonometric Integrals

<https://cheatography.com/crossant/cheat-sheets/integral-trigonometry/>

Integration by Partial Fractions

$(px+q)/(x-a)(x-b) = A/(x-a) + B/(x-b)$

$(px+q)/(x-a)^2 = A/(x-a) + B/(x-a)^2$

$(px^2+qx+r)/(x-a)(x-b)(x-c) = A/(x-a) + B/(x-b) + C/(x-c)$

$(px^2+qx+r)/(x-a)^2(x-b) = A/(x-a) + B/(x-a)^2 + C/(x-b)$

$(px^2+qx+r)/(x-a)(x^2+bx+c) = A/(x-a) + Bx+C/(x^2+bx+c)$

$\int 1/(a^2+x^2) dx = (1/a) \arctan(x/a) + C$

Integration by Partial Fractions is used to simplify integrals of polynomial rational expressions into simpler fractions with a factored, irreducible denominator

The degree (highest power) of the numerator's polynomial must be less than the degree of the denominator's polynomial, otherwise, polynomial long

Improper Integrals (cont)

Convergence of $\int f(x) dx$ $\lim_{t \rightarrow \pm\infty} f(x) dx = L$, where L is a constant

Divergence of $\int f(x) dx$ $\lim_{t \rightarrow \pm\infty} f(x) dx = \pm\infty$ or DNE

Improper Integrals are integrals with bounds at infinity (Type 1) or at least one discontinuity on the integrated region (Type 2)

Conic Sections

<https://cheatography.com/crossant/cheat-sheets/conic-sections/>

Parametric Curves and Polar Functions

| | |
|--|--|
| Function | $\int_a^b \sqrt{1+(f'(x))^2} dx$ |
| Parametric Function | $\int_a^b \sqrt{(x'(t))^2+(y'(t))^2} dt$ |
| Polar Function | $\int_a^b \sqrt{(r(\theta))^2+(r'(\theta))^2} d\theta$ |
| For standard functions: $f'(x)=-dy/dx$ | |
| For parametric functions: $x'(t)=-dx/dt$ and $y'(t)=dy/dt$ | |
| For polar functions: $r'(\theta)=dr/d\theta$ | |

integrals"

Choosing the "dv" term depends on what will simplify the integral the best, while being relatively simple to integrate

The constant of integration does not need to be inserted until the integral has been fully simplified

division must be used before converting the expression into partial fractions

Improper Integrals

$$\int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

$$\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

$$\int_{-\infty}^\infty f(x) dx = \lim_{t \rightarrow -\infty} \int_t^c f(x) dx + \lim_{t \rightarrow \infty} \int_c^t f(x) dx$$

| | |
|--|--|
| Parametric Curve C as a function of Parameter t | $(x,y)=(f(t),g(t))$ for t on [a,b] |
| Slope at a given point | $dy/dx = (dy/dt)/(dx/dt)$ |
| Second derivative | $d^2y/dx^2 = (dy/dt) / (dx/dt)^2$ |
| Polar Curve C as a function of Parameters r and θ | $(r,\theta)=(r,\theta \pm 2\pi n) = (-r,\theta \pm \pi n)$ |
| Slope at a given point | $dy/dx = (dy/d\theta)/(dx/d\theta)$ |
| Cartesian/Rectangular to Polar coordinates | $x=r\cos(\theta)$, $y=r\sin(\theta)$ |
| Polar to Cartesian/Rectangular coordinates | $r^2=x^2+y^2$ or $r=\sqrt{x^2+y^2}$, $\tan\theta=y/x$ or $\theta=\arctan(y/x)$ |
| $(dx/dt) \neq 0, (dx/d\theta) \neq 0$ | |



By **CROSSANT** (CROSSANT)
cheatography.com/crossant/

Published 28th May, 2023.
 Last updated 10th April, 2025.
 Page 4 of 5.

Sponsored by **Readable.com**
 Measure your website readability!
<https://readable.com>

Integral Approximations and Error Bounds

Midpoint Rule $\Delta x(f(\bar{x}_1)+f(\bar{x}_2)+f(\bar{x}_3)+\dots+f(\bar{x}_{n-1})+f(\bar{x}_n))$

Trapezoidal Rule $(\Delta x/2)(f(x_1)+2f(x_2)+2f(x_3)+\dots+2f(x_{n-1})+f(x_n))$

Simpson's Rule $(\Delta x/3)(f(x_1)+4f(x_2)+2f(x_3)+4f(x_4)+2f(x_5)+\dots+2f(x_{n-2})+4f(x_{n-1})+f(x_n))$

Midpoint Rule Error Bound $|E_m| \leq k(b-a)^3/24n^2$, $k=f''(x)_{\max}$ on $[a,b]$

Trapezoidal Rule Error Bound $|E_t| \leq k(b-a)^3/12n^2$, $k=f''(x)_{\max}$ on $[a,b]$

Simpson's Rule Error Bound $|E_s| \leq k(b-a)^5/180n^4$, $k=f^{(4)}(x)_{\max}$ on $[a,b]$

Integral Approximations are typically used to evaluate an integral that is very difficult or impossible to integrate

$$\Delta x = (b-a)/n$$

$\bar{x} = (x_{i-1} + x_i)/2$, the average/midpoint of two points x_{i-1} and x_i

Simpson's Rule can only be used if the given n is even, that is, $n=2k$ for some integer k

In order of most accurate to least accurate approximation:
Simpson's Rule, Midpoint Rule, Trapezoidal Rule, Left/Right endpoint approximation

