

### Lone Cases

Case Name	Step 1 - Exponent Manipulation	Step 2 - Trigonometric Identities	Step 3	Step 4
Sine-Odd	Manipulate to $\sin(x)(\sin^2(x))^p$	Sine Pythagorean Id.	Let $u=\cos(x)$	
Sine-Even	Manipulate to $(\sin^2(x))^p$	Sine Power-Reducing Id.	Expand	Solve cosine cases
Cosine-Odd	Manipulate to $\cos(x)(\cos^2(x))^p$	Cosine Pythagorean Id.	Let $u=\sin(x)$	
Cosine-Even	Manipulate to $(\cos^2(x))^p$	Cosine Power-Reducing Id.	Expand	Solve cosine cases
Tangent-Odd	Manipulate to $\tan(x)(\tan^2(x))^p$	Tangent Pythagorean Id.	Apply integral linearity	Solve other cases
Tangent-Even	Manipulate to $(\tan^2(x))^p$	Tangent Pythagorean Id.	Expand	Solve secant cases
Cosecant-Odd	Manipulate to $\csc(x)(\csc^2(x))^p$	Cosecant Pythagorean Id.	Expand and apply integral linearity	Solve other cases
Cosecant-Even	Manipulate to $\csc^2(x)(\csc^2(x))^p$	Cosecant Pythagorean Id.	Let $u=\cot(x)$	
Secant-Odd	Manipulate to $\sec(x)(\sec^2(x))^p$	Secant Pythagorean Id.	Expand and apply integral linearity	Solve other cases
Secant-Even	Manipulate to $\sec^2(x)(\sec^2(x))^p$	Secant Pythagorean Id.	Let $u=\tan(x)$	
Cotangent-Odd	Manipulate to $\cot(x)(\cot^2(x))^p$	Cotangent Pythagorean Id.	Apply integral linearity	Solve other cases
Cotangent-Even	Manipulate to $(\cot^2(x))^p$	Cotangent Pythagorean Id.	Expand	Solve cosecant cases

### Pair Cases

Case Name	Step 1 - Exponent Manipulation	Step 2	Step 3	Step 4
Sine-Odd and Cosine-Even	Manipulate sine powers to $\sin(x)(\sin^2(x))^p$	Sine Pythagorean Id.	Let $u=\cos(x)$	
Sine-Even and Cosine-Odd	Manipulate sine powers to $\cos(x)(\cos^2(x))^p$	Cosine Pythagorean Id.	Let $u=\sin(x)$	
Sine-Odd and Cosine-Odd	<i>Either two previous cases work</i>			
Sine-Even and Cosine-Even	Manipulate sine powers to $(\sin^2(x))^p$	Sine Pythagorean Id.	Expand and apply integral linearity	Solve cosine cases



By CROSSANT (CROSSANT)  
[cheatography.com/crossant/](https://cheatography.com/crossant/)

Published 6th July, 2024.  
 Last updated 6th July, 2024.  
 Page 1 of 3.

Sponsored by [ApolloPad.com](https://apollopad.com)  
 Everyone has a novel in them. Finish  
 Yours!  
<https://apollopad.com>

### Pair Cases (cont)

Sine-Cosine Same	Manipulate powers to $(\sin(x)\cos(x))^p$	Sine Double-Angle Id.	Solve sine cases	
Secant-Odd and Tangent-Even	Manipulate tangent powers to $(\tan^2(x))^p$	Tangent Pythagorean Id.	Expand and apply integral linearity	Solve secant cases
Secant-Even and Tangent-Odd	Manipulate powers to $\sec(x)\tan(x)\sec^p(x)\tan^q(x)$	Manipulate tangent powers to $(\tan^2(x))^r$	Tangent Pythagorean Id.	Let $u=\sec(x)$
Secant-Odd and Tangent-Odd	<i>Same as previous case</i>			
Secant-Even and Tangent-Even	Manipulate secant powers to $\sec^2(x)(\sec^2(x))^p$	Secant Pythagorean Id.	Let $u=\tan(x)$	
Secant-Tangent Same	<i>Same as previous two cases</i>			
Cosecant-Odd and Cotangent-Even	Manipulate cotangent powers to $(\cot^2(x))^p$	Cotangent Pythagorean Id.	Expand and apply integral linearity	Solve cosecant cases
Cosecant-Even and Cotangent-Odd	Manipulate powers to $\csc(x)\cot(x)\csc^p(x)\cot^q(x)$	Manipulate cotangent powers to $(\cot^2(x))^r$	Cotangent Pythagorean Id.	Let $u=\csc(x)$
Cosecant-Odd and Cotangent-Odd	<i>Same as previous case</i>			
Cosecant-Even and Cotangent-Even	Manipulate cosecant powers to $\csc^2(x)(\csc^2(x))^p$	Cosecant Pythagorean Id.	Let $u=\cot(x)$	
Cosecant-Cotangent Same	<i>Same as previous two cases</i>			

## Pythagorean Identities

Sine Pythagorean	$\sin^2(x) = 1 - \cos^2(x)$
Cosine Pythagorean	$\cos^2(x) = 1 - \sin^2(x)$
Tangent Pythagorean	$\tan^2(x) = \sec^2(x) - 1$
Cosecant Pythagorean	$\csc^2(x) = \cot^2(x) + 1$
Secant Pythagorean	$\sec^2(x) = \tan^2(x) + 1$
Cotangent Pythagorean	$\cot^2(x) = \csc^2(x) - 1$
Sine-Cosine Pythagorean	$1 = \sin^2(x) + \cos^2(x)$
Secant-Tangent Pythagorean	$1 = \sec^2(x) - \tan^2(x)$
Cosecant-Cotangent Pythagorean	$1 = \csc^2(x) - \cot^2(x)$

## Miscellaneous Information

Binomial Expansion	$(a+b)^n = \sum_{k=0}^n {}^n_k a^{n-k} b^k$
Integral Linearity Property: Sum and Difference	$\int (f(x) \pm g(x)) dx = \int f(x) dx \pm \int g(x) dx$
$\int \tan(x) dx$	$\ln \sec(x)  + C$
$\int \cot(x) dx$	$\ln \sin(x)  + C$
$\int \sec(x) dx$	$\ln \sec(x) + \tan(x)  + C$
$\int \sec^3(x) dx$	$\frac{1}{2}(\sec(x)\tan(x) + \ln \sec(x) + \tan(x) ) + C$
$\int \csc(x) dx$	$-\ln \csc(x) + \cot(x)  + C$
$\int \csc^3(x) dx$	$-\frac{1}{2}(\csc(x)\cot(x) + \ln \csc(x) + \cot(x) ) + C$

${}^n_k$  are the binomial coefficients, equal to  $n!/(k!(n-k)!)$



By CROSSANT (CROSSANT)  
[cheatography.com/crossant/](https://cheatography.com/crossant/)

Published 6th July, 2024.  
Last updated 6th July, 2024.  
Page 3 of 3.

Sponsored by [ApolloPad.com](https://apollopad.com)  
Everyone has a novel in them. Finish  
Yours!  
<https://apollopad.com>