

Circular Functions Definitions

Name	Right-Triangle Definition	Domain	Range
Sine Function	$\sin(\theta)=o/h$	$(-\infty, \infty)$	$[-1, 1]$
Cosine Function	$\cos(\theta)=a/h$	$(-\infty, \infty)$	$[-1, 1]$
Tangent Function	$\tan(\theta)=o/a$	$\{\theta \theta \neq \pi/2 + k\pi\}$	$(-\infty, \infty)$
Cosecant Function	$\csc(\theta)=h/o$	$\{\theta \theta \neq \pm k\pi\}$	$(-\infty, -1] \cup [1, \infty)$
Secant Function	$\sec(\theta)=h/a$	$\{\theta \theta \neq \pi/2 + k\pi\}$	$(-\infty, -1] \cup [1, \infty)$
Cotangent Function	$\cot(\theta)=a/o$	$\{\theta \theta \neq \pm k\pi\}$	$(-\infty, \infty)$
Inverse Sine Function	$\arcsin(o/h)=\theta$	$[-1, 1]$	$[-\pi/2, \pi/2]$
Inverse Cosine Function	$\arccos(a/h)=\theta$	$[-1, 1]$	$[0, \pi]$
Inverse Tangent Function	$\arctan(o/a)=\theta$	$(-\infty, \infty)$	$(-1, 1)$
Inverse Cosecant Function	$\text{arccsc}(h/o)=\theta$	$(-\infty, -1) \cup (1, \infty)$	$[-\pi/2, 0) \cup (0, \pi/2]$
Inverse Secant Function	$\text{arcsec}(h/a)=\theta$	$(-\infty, -1) \cup (1, \infty)$	$[0, \pi/2) \cup (\pi/2, \pi]$
Inverse Cotangent Function	$\text{arccot}(a/o)=\theta$	$(-\infty, \infty)$	$(0, 1)$
Circular Euler Relation	$e^{\pm i\theta}=\cos(\theta)\pm i\sin(\theta)$		
De Moivre's Theorem	$e^{in\theta}=(\cos(\theta)+i\sin(\theta))^n=\cos(n\theta)+i\sin(n\theta)$		

$n \in \mathbb{N}_1 = \{1, 2, 3, 4, 5, \dots\}$

"h" is the "hypotenuse" leg of a right triangle. It is directly across the right (90°) angle, and it has the longest length of the three sides.

"o" is the "opposite" leg of a right triangle. It is directly across the chosen angle θ .

"a" is the "adjacent" leg of a right triangle. It is the leg that is neither the hypotenuse leg, nor the opposite leg.

By the Pythagorean theorem, $o^2+a^2=h^2$

Hyperbolic Functions Definitions

Name	Exponential Definition	Domain	Range
Hyperbolic Sine Function	$\sinh(\theta)=(e^\theta-e^{-\theta})/2$	$(-\infty, \infty)$	$(-\infty, \infty)$
Hyperbolic Cosine Function	$\cosh(\theta)=(e^\theta+e^{-\theta})/2$	$(-\infty, \infty)$	$[1, \infty)$
Hyperbolic Tangent Function	$\tanh(\theta)=(e^\theta-e^{-\theta})/(e^\theta+e^{-\theta})$	$(-\infty, \infty)$	$(-1, 1)$
Hyperbolic Cosecant Function	$\text{csch}(\theta)=2/(e^\theta-e^{-\theta})$	$(-\infty, 0) \cup (0, \infty)$	$(-\infty, 0) \cup (0, \infty)$
Hyperbolic Secant Function	$\text{sech}(\theta)=2/(e^\theta+e^{-\theta})$	$(-\infty, \infty)$	$(0, 1]$



By CROSSANT (CROSSANT)
cheatography.com/crossant/

Published 26th August, 2024.
 Last updated 30th May, 2025.
 Page 1 of 9.

Sponsored by [CrosswordCheats.com](http://crosswordcheats.com)
 Learn to solve cryptic crosswords!
<http://crosswordcheats.com>

Hyperbolic Functions Definitions (cont)

Hyperbolic Cotangent Function	$\coth(\theta) = (e^\theta + e^{-\theta}) / (e^\theta - e^{-\theta})$	$(-\infty, 0) \cup (0, \infty)$	$(-\infty, -1) \cup (1, \infty)$
Inverse Hyperbolic Sine Function	$\operatorname{arcsinh}(x) = \ln(x + \sqrt{x^2 + 1})$	$(-\infty, \infty)$	$(-\infty, \infty)$
Inverse Hyperbolic Cosine Function	$\operatorname{arccosh}(x) = \ln(x + \sqrt{x^2 - 1})$	$[1, \infty)$	$[0, \infty)$
Inverse Hyperbolic Tangent Function	$\operatorname{arctanh}(x) = \frac{1}{2} \ln((1+x)/(1-x))$	$(-1, 1)$	$(-\infty, \infty)$
Inverse Hyperbolic Cosecant Function	$\operatorname{arccsch}(x) = \ln((1 \pm \sqrt{1+x^2})/x)$	$(-\infty, 0) \cup (0, \infty)$	$(-\infty, 0) \cup (0, \infty)$
Inverse Hyperbolic Secant Function	$\operatorname{arcsech}(x) = \ln((1 + \sqrt{1-x^2})/x)$	$(0, 1]$	$[0, \infty)$
Inverse Hyperbolic Cotangent Function	$\operatorname{arccoth}(x) = \frac{1}{2} \ln((x+1)/(x-1))$	$(-\infty, -1) \cup (1, \infty)$	$(-\infty, 0) \cup (0, \infty)$
Hyperbolic Euler Relation	$e^{\pm\theta} = \cosh(\theta) \pm \sinh(\theta)$		
De Moivre's Theorem (Hyperbolic)	$e^{n\theta} = (\cosh(\theta) + \sinh(\theta))^n = \cosh(n\theta) + \sinh(n\theta)$		
$n \in \mathbb{N}_1 = \{1, 2, 3, 4, 5, \dots\}$			

Complex Definitions

Name	Complex Relation	Circular-Hyperbolic Relation
Complex Sine	$\sin(z) = (e^{iz} - e^{-iz})/2i$	$\sin(z) = i \sinh(iz)$
Complex Cosine	$\cos(z) = (e^{iz} + e^{-iz})/2$	$\cos(z) = \cosh(iz)$
Complex Tangent	$\tan(z) = -i(e^{iz} - e^{-iz})/(e^{iz} + e^{-iz})$	$\tan(z) = -i \tanh(iz)$
Complex Cosecant	$\csc(z) = 2i/(e^{iz} - e^{-iz})$	$\csc(z) = i \operatorname{csch}(iz)$
Complex Secant	$\sec(z) = 2/(e^{iz} + e^{-iz})$	$\sec(z) = i \operatorname{sech}(iz)$
Complex Cotangent	$\cot(z) = i(e^{iz} + e^{-iz})/(e^{iz} - e^{-iz})$	$\cot(z) = i \operatorname{coth}(iz)$
Complex Inverse Sine	$\operatorname{arcsin}(z) = -i \ln(iz \pm \sqrt{(1-z^2)})$	$\operatorname{arcsin}(z) = i \operatorname{arsinh}(iz)$
Complex Inverse Cosine	$\operatorname{arccos}(z) = -i \ln(z \pm i\sqrt{(1-z^2)})$	$\operatorname{arccos}(z) = \pm i \operatorname{arcosh}(z)$
Complex Inverse Tangent	$\operatorname{arctan}(z) = (i/2) \ln((i+z)/(i-z))$	$\operatorname{arctan}(z) = i \operatorname{arctanh}(iz)$
Complex Inverse Cosecant	$\operatorname{arccsc}(z) = -i \ln((i+\sqrt{z^2-1})/z)$	$\operatorname{arccsc}(z) = i \operatorname{arccsch}(iz)$
Complex Inverse Secant	$\operatorname{arcsec}(z) = -i \ln((1+\sqrt{1-z^2})/z)$	$\operatorname{arcsec}(z) = \pm i \operatorname{arcsech}(z)$
Complex Inverse Cotangent	$\operatorname{arccot}(z) = -(i/2) \ln((z+i)/(z-i))$	$\operatorname{arccot}(z) = \pm i \operatorname{arccoth}(iz)$
Complex Hyperbolic Sine	None	$\sinh(z) = -i \sin(iz)$



Complex Definitions (cont)

Complex Hyperbolic Cosine	None	$\cosh(z)=\cos(iz)$
Complex Hyperbolic Tangent	None	$\tanh(z)=-i\tan(iz)$
Complex Hyperbolic Cosecant	None	$\operatorname{csch}(z)=i\csc(iz)$
Complex Hyperbolic Secant	None	$\operatorname{sech}(z)=\sec(iz)$
Complex Hyperbolic Cotangent	None	$\coth(z)=i\cot(iz)$
Complex Inverse Hyperbolic Sine	None	$\operatorname{arsinh}(z)=-i\operatorname{arcsin}(iz)$
Complex Inverse Hyperbolic Cosine	None	$\operatorname{arcosh}(z)=\pm i\operatorname{arccos}(z)$
Complex Inverse Hyperbolic Tangent	None	$\operatorname{artanh}(z)=-i\operatorname{arctan}(iz)$
Complex Inverse Hyperbolic Cosecant	None	$\operatorname{arccsch}(z)=-i\operatorname{arccsc}(iz)$
Complex Inverse Hyperbolic Secant	None	$\operatorname{arcsech}(z)=\pm i\operatorname{arcsec}(z)$
Complex Inverse Hyperbolic Cotangent	None	$\operatorname{arccoth}(z)=-i\operatorname{arccot}(iz)$

$i=\sqrt{-1}$

z is a complex variable of the form $a+bi$, where a and b are real numbers, and i is the imaginary unit

Circular Functions Unit Circle Values

θ (Radians)	θ (Degrees)	$\sin(\theta)$	$\cos(\theta)$	$\tan(\theta)$	$\csc(\theta)$	$\sec(\theta)$	$\cot(\theta)$
0	0°	0	1	0	undefined	1	undefined
$\pi/6$	30°	1/2	$\sqrt{3}/2$	$\sqrt{3}/3$	2	$2\sqrt{3}/3$	$\sqrt{3}$
$\pi/4$	45°	$\sqrt{2}/2$	$\sqrt{2}/2$	1	$\sqrt{2}$	$\sqrt{2}$	1
$\pi/3$	60°	$\sqrt{3}/2$	1/2	$\sqrt{3}$	$2\sqrt{3}/3$	2	$\sqrt{3}/3$
$\pi/2$	90°	1	0	undefined	1	undefined	0
$2\pi/3$	120°	$-\sqrt{3}/2$	-1/2	- $\sqrt{3}$	$2\sqrt{3}/3$	-2	$-\sqrt{3}/3$
$3\pi/4$	135°	$-\sqrt{2}/2$	$-\sqrt{2}/2$	-1	$\sqrt{2}$	$-\sqrt{2}$	-1
$5\pi/6$	150°	1/2	$-\sqrt{3}/2$	$-\sqrt{3}/3$	2	$-2\sqrt{3}/3$	$-\sqrt{3}$
π	180°	0	-1	0	undefined	-1	undefined
$7\pi/6$	210°	-1/2	$-\sqrt{3}/2$	$\sqrt{3}/3$	-2	$-2\sqrt{3}/3$	$\sqrt{3}$
$5\pi/4$	225°	$-\sqrt{2}/2$	$-\sqrt{2}/2$	1	$-\sqrt{2}$	$-\sqrt{2}$	1
$4\pi/3$	240°	$-\sqrt{3}/2$	-1/2	$\sqrt{3}$	$-2\sqrt{3}/3$	-2	$\sqrt{3}/3$



By CROSSANT (CROSSANT)
cheatography.com/crossant/

Published 26th August, 2024.
 Last updated 30th May, 2025.
 Page 3 of 9.

Sponsored by [CrosswordCheats.com](http://crosswordcheats.com)
 Learn to solve cryptic crosswords!
<http://crosswordcheats.com>

Circular Functions Unit Circle Values (cont)

$3\pi/2$	270°	-1	0	undefined	-1	undefined	0
$5\pi/3$	300°	$-\sqrt{3}/2$	$1/2$	$-\sqrt{3}$	$-2\sqrt{3}/3$	2	$-\sqrt{3}/3$
$7\pi/4$	315°	$-\sqrt{2}/2$	$\sqrt{2}/2$	-1	$-\sqrt{2}$	$\sqrt{2}$	-1
$11\pi/6$	330°	$-1/2$	$\sqrt{3}/2$	$-\sqrt{3}/3$	-2	$2\sqrt{3}/3$	$-\sqrt{3}$
2π	360°	0	1	0	undefined	1	undefined

The coordinates $(\cos(\theta), \sin(\theta))$ represent x and y coordinates of θ on the unit circle $x^2+y^2=1$

Circular Compositional Identities

Composition	$\sin(x)$	$\cos(x)$	$\tan(x)$
$\arcsin(x)$	x	$\sqrt{1-x^2}$	$x/\sqrt{1-x^2}$
$\arccos(x)$	$\sqrt{1-x^2}$	x	$\sqrt{1-x^2}/x$
$\arctan(x)$	$x/\sqrt{1+x^2}$	$1/\sqrt{1+x^2}$	x
$\text{arcsc}(x)$	$1/x$	$\sqrt{x^2-1}/ x $	$\pm 1/\sqrt{x^2-1}$
$\text{arcsec}(x)$	$\sqrt{x^2-1}/ x $	$1/x$	$\pm\sqrt{x^2-1}$
$\text{arccot}(x)$	$1/\sqrt{1+x^2}$	$x/\sqrt{1+x^2}$	$1/x$

Each composition is valid on different domains

Hyperbolic Compositional Identities

Composition	$\sinh(x)$	$\cosh(x)$	$\tanh(x)$
$\text{arcsinh}(x)$	x	$\sqrt{1+x^2}$	$x/\sqrt{1-x^2}$
$\text{arccosh}(x)$	$\sqrt{x^2-1}$	x	$\sqrt{x^2-1}/x$
$\text{arctanh}(x)$	$x/\sqrt{1-x^2}$	$1/\sqrt{1-x^2}$	x
$\text{arccsch}(x)$	$1/x$	$\sqrt{x^2+1}/ x $	$1/\sqrt{x^2+1}$
$\text{arcsech}(x)$	$\sqrt{1-x^2}/x$	$1/x$	$\sqrt{1-x^2}$
$\text{arccoth}(x)$	$x/\sqrt{1-x^2}$	$ x /\sqrt{x^2-1}$	$1/x$

Each composition is valid on different domains

Circular Quotient & Reciprocal Identities

Cofunctional Phase Shift Properties (cont)

Cosecant Complimentary	$\csc(\theta)=\sec(\pi/2-\theta)$
Cosecant Supplementary	$\csc(\theta)=\csc(\pi-\theta)$
Secant Complementary	$\sec(\theta)=\csc(\pi/2-\theta)$
Secant Supplementary	$\sec(\theta)=-\sec(\pi-\theta)$
Cotangent Complimentary	$\cot(\theta)=\tan(\pi/2-\theta)$
Cotangent Supplementary	$\cot(\theta)=-\cot(\pi-\theta)$

$$n \in \mathbb{N}_1 = \{1, 2, 3, 4, 5, \dots\}$$

Circular Parity Properties

Sine Odd	$\sin(-\theta)=-\sin(\theta)$
Cosine Even	$\cos(-\theta)=\cos(\theta)$
Tangent Odd	$\tan(-\theta)=-\tan(\theta)$
Cosecant Odd	$\csc(-\theta)=-\csc(\theta)$
Secant Even	$\sec(-\theta)=\sec(\theta)$
Cotangent Odd	$\cot(-\theta)=-\cot(\theta)$

(C) Half/Multiple-Angle Identities (cont)

Circular Pythagorean Identities

Periodicity Properties

Tangent Quotient	$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$	Sine Periodicity	$\sin(\theta) = \sin(\theta + 2\pi n)$	Sine-Cosine	$\sin^2(\theta) + \cos^2(\theta) = 1$	Tangent Half-Angle 2	$\tan(\theta/2) = \frac{1 - \cos(\theta)}{\sin(\theta)}$
Cotangent Quotient	$\cot(\theta) = \frac{\cos(\theta)}{\sin(\theta)}$	Cosine Periodicity	$\cos(\theta) = \cos(\theta + 2\pi n)$	Pythagorean		Tangent Half-Angle 3	$\tan(\theta/2) = \frac{\sin(\theta)}{(1 + \cos(\theta))}$
Sine Reciprocal	$\sin(\theta) = 1/\csc(\theta)$	Tangent Periodicity	$\tan(\theta) = \tan(\theta + \pi n)$	Secant-Tangent	$\tan^2(\theta) + 1 = \sec^2(\theta)$	Sine Double-Angle 1	$\sin(2\theta) = 2\sin(\theta)\cos(\theta)$
Cosine Reciprocal	$\cos(\theta) = 1/\sec(\theta)$	Cosecant Periodicity	$\csc(\theta) = \csc(\theta + 2\pi n)$	Pythagorean		Sine Double-Angle 2	$\sin(2\theta) = 2\tan(\theta)/(1 + \tan^2(\theta))$
Tangent Reciprocal	$\tan(\theta) = 1/\cot(\theta)$	Secant Periodicity	$\sec(\theta) = \sec(\theta + 2\pi n)$	Cosecant-Cotangent	$1 + \cot^2(\theta) = \csc^2(\theta)$	Cosine Double-Angle 1	$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta)$
Cosecant Reciprocal	$\csc(\theta) = 1/\sin(\theta)$	Cotangent Periodicity	$\cot(\theta) = \cot(\theta + \pi n)$	Pythagorean		Cosine Double-Angle 2	$\cos(2\theta) = 2\cos^2(\theta) - 1$
Secant Reciprocal	$\sec(\theta) = 1/\cos(\theta)$	$n \in \mathbb{N}_1 = \{1, 2, 3, 4, 5, \dots\}$		The last two Pythagorean Identities are obtained by dividing all the terms of the original Sine-Cosine Identity by $\cos^2(\theta)$ and $\sin^2(\theta)$, respectively		Cosine Double-Angle 3	$\cos(2\theta) = 1 - 2\sin^2(\theta)$
Cotangent Reciprocal	$\cot(\theta) = 1/\tan(\theta)$			(C) Half/Multiple-Angle Identities		Cosine Double-Angle 4	$\cos(2\theta) = (1 - \tan^2(\theta))/(-1 + \tan^2(\theta))$
All the following identities are true for values that do not cause division by zero		Sine Half-Angle	$\sin(\theta/2) = \pm \sqrt{\frac{1}{2}(1 - \cos(\theta))}$	Tangent Half-Angle 1	$\tan(\theta/2) = \pm \sqrt{\frac{(1 - \cos(\theta))}{(1 + \cos(\theta))}}$	Tangent Double-Angle 1	$\tan(2\theta) = 2\tan(\theta)/(1 - \tan^2(\theta))$
		Cosine Half-Angle	$\cos(\theta/2) = \pm \sqrt{\frac{1}{2}(1 + \cos(\theta))}$			Tangent Double-Angle 2	$\tan(2\theta) = 2/(\cot(\theta) - \tan(\theta))$
						Sine Triple-Angle	$\sin(3\theta) = 3\sin(\theta) - 4\sin^3(\theta)$



By CROSSANT (CROSSANT)
cheatography.com/crossant/


Published 26th August, 2024.
 Last updated 30th May, 2025.
 Page 5 of 9.

Sponsored by [CrosswordCheats.com](http://crosswordcheats.com)
 Learn to solve cryptic crosswords!
<http://crosswordcheats.com>

Cheatography

Trigonometric Properties and Identities Cheat Sheet

by CROSSANT (CROSSANT) via cheatography.com/186482/cs/39959/

(C) Half/Multiple-Angle Identities (cont)		Circular Sum/Difference/Product Identities (cont)		Circular-Inverse Reciprocal Identities		Circular-Inverse Negative Input Identities	
Cosine Triple-Angle	$\cos(3\theta)=4\cos^3(\theta)-3\cos(\theta)$	Cosine Product-Sum	$\cos(\theta)-\cos(\varphi) = \frac{1}{2}(\cos(-\theta-\varphi)+\cos(-\theta+\varphi))$	Sine Reciprocal	$\arcsin(1/x)=\arcsin(x)$	Sine Odd	$\arcsin(-x)=-\arcsin(x)$
Tangent Triple-Angle	$\tan(3\theta)=(3\tan(\theta)-\tan^3(\theta))/(1-3\tan^2(\theta))$	Sine-Cosine Product-Sum	$\sin(\theta)-\cos(\varphi) = \frac{1}{2}(\sin(-\theta-\varphi)+\sin(-\theta+\varphi))$	Cosine Reciprocal	$\arccos(1/x)=\arccos(x)$	Cosine Translation	$\arccos(-x)=\pi-\arccos(x)$
Sine Multiple-Angle Formula: $\sin(n\theta)=\sum_{k=0}^n (^n_k)\cos^k(\theta)\sin^{n-k}(\theta)\sin((\pi/2)(n-k))$		Tangent Sum/Difference	$\tan(\theta\pm\varphi) = \frac{(\tan(\theta)\pm\tan(-\varphi))/(1-\tan(\theta)\tan(\varphi))}{\tan(\varphi)}$	Tangent Reciprocal 1	$\arctan(1/x)=\arctan(x), x>0$	Tangent Odd	$\arctan(-x)=-\arctan(x)$
Cosine Multiple-Angle Formula: $\cos(n\theta)=\sum_{k=0}^n (^n_k)\cos^k(\theta)\sin^{n-k}(\theta)\cos((\pi/2)(n-k))$		Tangent Sum	$\tan(\theta\pm\varphi) = \frac{\sin(\theta\pm\varphi)/(\cos(\theta)\cos(\varphi))}{\pm\tan(\varphi)}$	Tangent Reciprocal 2	$\arctan(1/x)=\arctan(x), \pi<x<0$	Cosecant Odd	$\arccsc(-x)=-\arccsc(x)$
All the following identities are true for values that do not cause division by zero		Cosecant Reciprocal	$\arccsc(1/x)=\arcsin(x)$	Secant Reciprocal	$\arcsec(1/x)=\arccos(x)$	Secant Translation	$\arcsec(-x)=\pi-\arcsec(x)$
Circular Sum/Difference/Product Identities		Cotangent Reciprocal 1	$\arccot(1/x)=\arctan(x), x>0$	Cotangent Reciprocal 2	$\arccot(1/x)=\arctan(x)+\pi, x<0$	Cotangent Translation	$\arccot(-x)=\pi-\arccot(x)$
Sine Sum/Difference	$\sin(\theta\pm\varphi) = \sin(\theta)\cos(\varphi)\pm\cos(\theta)\sin(\varphi)$	Circular-Inverse Complimentary Identities		(CI) Half/Multiple Substitution Identities			
Sine Sum-Product	$\sin(\theta)\pm\sin(\varphi) = 2\sin((\theta\pm\varphi)/2)\cos((\theta\mp\varphi)/2)$	Tangent Product	$\tan(\theta)-\tan(\varphi) = (\tan(\theta)+\tan(-\varphi))/(\cot(\theta)+\cot(\varphi))$	Sine Complimentary	$\arcsin(x)=\pi/2-\arccos(x)$	Half Sine Substitution 1	$\frac{1}{2}\arcsin(x)=\arcsin(\sqrt{(1+x)/2})-\pi/4$
Sine Product-Sum	$\sin(\theta)\sin(\varphi) = \frac{1}{2}(\cos(-\theta-\varphi)-\cos(-\theta+\varphi))$	Tangent-Cotangent Product	$\tan(\theta)-\cot(\varphi) = (\tan(\theta)+\cot(-\varphi))/(\cot(\theta)+\tan(\varphi))$	Cosine Complimentary	$\arccos(x)=\pi/2-\arcsin(x)$	Half Sine Substitution 2	$\frac{1}{2}\arcsin(x)=\pi/4-\arcsin(\sqrt{(1-x)/2})$
Cosine Sum/Difference	$\cos(\theta\pm\varphi) = \cos(\theta)\cos(\varphi)-\sin(\theta)\sin(\varphi)$	Sine and Cosine Unit Circle		Tangent Complimentary	$\arctan(x)=\pi/2-2\arccot(x)$	Half Cosine Substitution 1	$\frac{1}{2}\arccos(x)=\arccos(\sqrt{(1+x)/2})$
Cosine Sum-Product	$\cos(\theta)\pm\cos(\varphi) = 2\cos((\theta\pm\varphi)/2)\cos((\theta\mp\varphi)/2)$	Secant Complimentary		Cosecant Complimentary	$\arccsc(x)=\pi/2-\arccsc(x)$	Half Cosine Substitution 2	$\frac{1}{2}\arccos(x)=\pi/2-\arccos(\sqrt{(1-x)/2})$
Cosine Product		Cotangent Complimentary	$\arccot(x)=\pi/2-\arctan(x)$	Double Sine Substitution			
$x^2+y^2=1$		Double Cosine Substitution 1		Double Sine Substitution	$2\arcsin(x)=\arcsin(2x\sqrt{1-x^2}), x /\leq\pi/2$	Double Cosine Substitution 1	
		Double Cosine Substitution 2		Double Cosine Substitution	$2\arccos(x)=\arccos(2x^2-1), x\geq 0$		



(CI) Half/Multiple Substitution Identities (cont)		Circular-Inverse Sum/Difference Identities		Law of Sines/Cosines/Tangents (cont)		Tangent Unit Circle
Double Cosine Substitution 2	$2\arccos(x)=2\pi-a-\arccos(2x^2-1), x \leq 0$	Sine Sum/Difference	$\arcsin(x) \pm \arcsin(y) = \arcsin(x\sqrt{1-y^2}) \pm y\sqrt{1-x^2}$	Law of Tangents 2	$(b-c)/(b+c) = \tan((\beta-\gamma)/2)/\tan((\beta+\gamma)/2)$	
Double Tangent Substitution 1	$2\arctan(x) = \arcsin(2x/(1+x^2)), x \leq 1$	Cosine Sum/Difference	$\arccos(x) \pm \arccos(y) = \arccos(xy \mp \sqrt{(1-x^2)\sqrt{(1-y^2)}})$	Law of Tangents 3	$(c-a)/(c+a) = \tan((\gamma-\alpha)/2)/\tan((\gamma+\alpha)/2)$	
Double Tangent Substitution 2	$2\arctan(x) = \pm \arccos((1-x^2)/(1+x^2))$	Cosine-Sine Sum/Difference	$\arccos(x) \pm \arcsin(y) = \arccos(x\sqrt{1-y^2}) \mp y\sqrt{1-x^2}$	Side lengths a, b, and c are opposite of the angles α, β, and γ, respectively.		
Double Tangent Substitution 3	$2\arctan(x) = \arctan(2x/(1-x^2)), x < 1$	Tangent Sum/Difference	$\arctan(x) \pm \arctan(y) = \arctan((x \pm y)/(1 \mp xy)), 1 \mp xy \neq 0$	Measurements And Formulas		
		Law of Sines/Cosines/Tangents		Radians-Degrees	1 radian = $180/\pi$ degrees; $1 = (180/\pi)^\circ$	
Triple Sine Substitution 1	$3\arcsin(x) = \arcsin(3x-4x^3), x \leq \frac{1}{2}$	Law of Sines 1	$\sin(\alpha)/a = \sin(\beta)/b = \sin(\gamma)/c$	Degrees-Radians	1 degree = $\pi/180$ radians; $1^\circ = \pi/180$ radians	
Triple Sine Substitution 2	$3\arcsin(x) = \arcsin(4x^3-3x) \pm \pi, x \geq \frac{1}{2}$	Law of Sines 2	$a/\sin(\alpha) = b/\sin(\beta) = c/\sin(\gamma)$	Degrees, Minutes, and Seconds	1 degree = 60 minutes = 3600 seconds; $1^\circ = 60' = 3600''$ (DMS)	
Triple Cosine Substitution 1	$3\arccos(x) = \arccos(3x-4x^3), x \leq \frac{1}{2}$	Law of Cosines 1	$a^2 = b^2 + c^2 - 2bc \cos(\alpha)$	Arc Length/Angular Displacement	$s = r\theta$ units	
Triple Cosine Substitution 2	$3\arccos(x) = \arccos(4x^3-3x) + \pi \pm \pi, x \geq \frac{1}{2}$	Law of Cosines 2	$b^2 = a^2 + c^2 - 2ac \cos(\beta)$	Sector Area	$\frac{1}{2}r^2\theta$ units ²	
Triple Tangent Substitution 1	$3\arctan(x) = \arctan((3x-x^3)/(1-3x^2)), x \leq \sqrt{3}/3$	Law of Cosines 3	$c^2 = a^2 + b^2 - 2ab \cos(\gamma)$	Area of a Triangle	$A_T = \frac{1}{2}bh$ units ²	
Triple Tangent Substitution 2	$3\arctan(x) = \arctan((3x-x^3)/(1-3x^2)) \pm \pi, x \geq \sqrt{3}/3$	Law of Tangents 1	$(a-b)/(a+b) = \tan((\alpha-\beta)/2)/\tan((\alpha+\beta)/2)$	Area of a Circle	$A_C = \pi r^2$ units ²	
				Pythagorean Theorem	$a^2 + b^2 = c^2$	All the following identities are true for values that do not cause division by zero
				Radians are unitless a, b, and c are side lengths of a right-triangle		



Hyperbolic Parity Properties		(H) Half-Angle & Multiple-Angle Identities (cont)		(H) Half-Angle & Multiple-Angle Identities (cont)		(H) Sum/Difference/Product Identities (cont)	
Sine Odd	$\sinh(-\theta) = -\sinh(\theta)$	Tangent	$\tanh(\theta/2) = \pm \sqrt{((\cosh(\theta)-1)/(\cosh(\theta)+1))}$	Sine	$\sinh(3\theta) = 3\sinh(\theta) + 4\sinh^3(\theta)$	Cosine	$\cosh(\theta) - \cosh(\varphi) = 2\sinh((\theta-\varphi)/2)\sinh((\theta+\varphi)/2)$
Cosine Even	$\cosh(-\theta) = \cosh(\theta)$	Half-Angle 1		Triple-Angle		Sum-Product 2	
Tangent Odd	$\tanh(-\theta) = -\tanh(\theta)$	Tangent	$\tanh(\theta/2) = (\cosh(\theta)-1)/\sinh(\theta)$	Cosine	$\cosh(3\theta) = 4\cosh^3(\theta) - 3\cosh(\theta)$	Cosine	$\cosh(\theta)\cosh(\varphi) = \frac{1}{2}(\cosh(\theta+\varphi) + \cosh(\theta-\varphi))$
Cosecant Odd	$\operatorname{csch}(-\theta) = -\operatorname{csch}(\theta)$	Angle 2		Tangent	$\tanh(3\theta) = (3\tan(\theta) + \tan^3(\theta))/(-1 + 3\tan^2(\theta))$	Sine-Cosine Product-Sum	$\sinh(\theta)\cosh(\varphi) = \frac{1}{2}(\sinh(\theta+\varphi) + \sinh(\theta-\varphi))$
Secant Even	$\operatorname{sech}(-\theta) = \operatorname{sech}(\theta)$	Tangent	$\tanh(\theta/2) = \sinh(\theta)/(\cosh(\theta)+1)$	Triple-Angle			
Cotangent Odd	$\coth(-\theta) = -\coth(\theta)$	Angle 3					
Hyperbolic Pythagorean Identities		Sine Double-Angle 1	$\sinh(2\theta) = 2\sinh(\theta)\cosh(\theta)$	(H) Sum/Difference/Product Identities		Tangent Sum-Difference	$\tanh(\theta \pm \varphi) = (\tanh(\theta) \pm \tanh(\varphi)) / (1 \pm \tanh(\theta)\tanh(\varphi))$
Sine-Cosine Pythagorean	$\cosh^2(\theta) - \sinh^2(\theta) = 1$	Sine Double-Angle 2	$\sinh(2\theta) = 2\tanh(\theta) / (1 - \tanh^2(\theta))$	Sine	$\sinh(\theta \pm \varphi)$	$\cosh(\theta \pm \varphi) = \cosh(\theta)\cosh(\varphi) \pm \sinh(\theta)\sinh(\varphi)$	
Secant Pythagorean	$1 - \tanh^2(\theta) = \operatorname{sech}^2(\theta)$	Cosine Double-Angle 1	$\cosh(2\theta) = \cosh^2(\theta) + \sinh^2(\theta)$	Sine	$\sinh(\theta) \pm \sinh(\varphi)$	$2\sinh((\theta \pm \varphi)/2)\cosh((\theta \mp \varphi)/2)$	
Cosecant Pythagorean	$\coth^2(\theta) - 1 = \operatorname{csch}^2(\theta)$	Cosine Double-Angle 2	$\cosh(2\theta) = 2\cosh^2(\theta) - 1$	Sine	$\sinh(\theta)\sinh(\varphi)$	$\frac{1}{2}(\cosh(\theta+\varphi) - \cosh(\theta-\varphi))$	
The last two Hyperbolic Pythagorean Identities are obtained by dividing all the terms of the original Sine-Cosine Identity by $\cosh^2(\theta)$ and $\sinh^2(\theta)$, respectively		Cosine Double-Angle 3	$\cosh(2\theta) = -1 + 2\sinh^2(\theta)$	Cosine	$\cosh(\theta \pm \varphi)$	$\cosh(\theta \pm \varphi) = \cosh(\theta)\cosh(\varphi) \mp \sinh(\theta)\sinh(\varphi)$	
		Cosine Double-Angle 4	$\cosh(2\theta) = (-1 + \tanh^2(\theta)) / (1 - \tanh^2(\theta))$	Tangent	$\tanh(\theta \pm \varphi)$	$\tanh(\theta \pm \varphi) = (\tanh(\theta) \pm \tanh(\varphi)) / (1 \pm \tanh(\theta)\tanh(\varphi))$	
		Tangent Double-Angle 1	$\tanh(2\theta) = 2\tanh(\theta) / (1 + \tanh^2(\theta))$	Cosine	$\cosh(\theta) + \cosh(\varphi)$	$2\cosh((\theta + \varphi)/2)\cosh((\theta - \varphi)/2)$	
		Tangent Double-Angle 2	$\tanh(2\theta) = 2/(\cosh(\theta) + \tanh(\theta))$	Tangent-Cotangent Product	$\tanh(\theta)\cot(\varphi)$	$(\tanh(\theta) + \tanh(\varphi)) / (\cosh(\theta) + \cosh(\varphi))$	
(H) Half-Angle & Multiple-Angle Identities							
Sine Half-Angle	$\sinh(\theta/2) = \pm \sqrt{(\frac{1}{2}(\cosh(\theta)-1))}$						
Cosine Half-Angle	$\cosh(\theta/2) = \sqrt{(\frac{1}{2}(\cosh(\theta)+1))}$						



Right-Triangle Relations

$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$	$\csc \theta = \frac{\text{hypotenuse}}{\text{opposite}}$
$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$	$\sec \theta = \frac{\text{hypotenuse}}{\text{adjacent}}$
$\tan \theta = \frac{\text{opposite}}{\text{adjacent}}$	$\cot \theta = \frac{\text{adjacent}}{\text{opposite}}$

Hyperbolic-Inverse Reciprocal Identities

Sine Reciprocal	$\operatorname{arcsinh}(1/x) = \operatorname{arccsch}(x)$
Cosine Reciprocal	$\operatorname{arccosh}(1/x) = \operatorname{arcsech}(x)$
Tangent Reciprocal	$\operatorname{arctanh}(1/x) = \operatorname{arccoth}(x)$
Cosecant Reciprocal	$\operatorname{arccsch}(1/x) = \operatorname{arcsinh}(x)$
Secant Reciprocal	$\operatorname{arcsech}(1/x) = \operatorname{arccosh}(x)$
Cotangent Reciprocal	$\operatorname{arccoth}(1/x) = \operatorname{arctanh}(x)$

(HI) Negative Input Identities

Inverse Sine Odd	$\operatorname{arcsinh}(-x) = -\operatorname{arcsinh}(x)$
Inverse Tangent Odd	$\operatorname{arctanh}(-x) = -\operatorname{arctanh}(x)$
Inverse Cosecant Odd	$\operatorname{arccsch}(-x) = -\operatorname{arccsch}(x)$
Inverse Cotangent Odd	$\operatorname{arccoth}(-x) = -\operatorname{arccoth}(x)$

(HI) Half/Multiple Substitution Identities

Half Sine Substitution	$\frac{1}{2}\operatorname{arcsinh}(x) = \pm \operatorname{arcsinh}(\sqrt{(\sqrt{(1+x^2)-1})/2})$
Half Cosine Substitution	$\frac{1}{2}\operatorname{arccosh}(x) = \operatorname{arccosh}(\sqrt{((x+1)/2)})$

Half Tangent Substitution	$\frac{1}{2}\operatorname{arctanh}(x) = \operatorname{arctanh}(x/(1+\sqrt{(1-x^2)}))$
---------------------------	---

Double Sine Substitution 1	$2\operatorname{arcsinh}(x) = \operatorname{arcsinh}(2x\sqrt{(1+x^2)})$
Double Sine Substitution 2	$2\operatorname{arcsinh}(x) = \pm \operatorname{arccosh}(2x^2+1)$

Double Cosine Substitution	$2\operatorname{arccosh}(x) = \operatorname{arccosh}(2x^2-1), x \geq 1$
Double Tangent Substitution	$2\operatorname{arctanh}(x) = \operatorname{arctanh}(2x/(1+x^2)), x < 1$

Triple Sine Substitution	$3\operatorname{arcsinh}(x) = \operatorname{arcsinh}(3x+4x^3)$
Triple Cosine Substitution	$3\operatorname{arccosh}(x) = \operatorname{arccosh}(4x^3-3x)$

Triple Tangent Substitution	$3\operatorname{arctanh}(x) = \operatorname{arctanh}((3x+x^3)/(1+3x^2))$
-----------------------------	--

(HI) Sum/Difference Identities

Sine Sum/Difference	$\operatorname{arcsinh}(x) \pm \operatorname{arcsinh}(y) = \operatorname{arcsinh}(x\sqrt{(y^2+1)} \pm y\sqrt{(x^2+1)})$
Cosine Sum/Difference	$\operatorname{arccosh}(x) \pm \operatorname{arccosh}(y) = \operatorname{arccosh}(xy \pm \sqrt{((x^2-1)(y^2-1))})$

Sine-Cosine	$\operatorname{arcsinh}(x) \pm \operatorname{arccosh}(y) = \operatorname{arcsinh}(xy \pm \sqrt{((x^2+1)(y^2-1))})$
-------------	--

Sine-Difference	$\operatorname{arctanh}(1) = \operatorname{arctanh}(x/(1+\sqrt{(1-x^2)}))$
-----------------	--

Sine-Cosine	$\operatorname{arccosh}(x) \pm \operatorname{arccosh}(y) = \operatorname{arccosh}(xy \pm \sqrt{((x^2-1)(y^2-1))})$
-------------	--

Tangent Sum/Difference	$\operatorname{arctanh}(x) \pm \operatorname{arctanh}(y) = \operatorname{arctanh}((x \pm y)/(1 \pm xy))$
------------------------	--

(HI) Logarithmic/Compositional Conversions

Sine Logarithmic	$\ln(x) = \operatorname{arcsinh}((x^2-1)/2x), x > 0$
------------------	--

Cosine Logarithmic	$\ln(x) = \pm \operatorname{arccosh}((x^2+1)/2x)$
--------------------	---

Tangent Logarithmic	$\ln(x) = \operatorname{arctanh}((x^2-1)/(x^2+1))$
---------------------	--

Sine Hyperbolic	$\operatorname{arcsinh}(\tan(x)) = \ln(\sec(x+\pi n)+\tan(x+\pi n))$
-----------------	--

Sine Circular	$\operatorname{arcsinh}(\tan(x)) = \pm \operatorname{arccosh}(\sec(x+\pi n))$
---------------	---

(HI) Logarithmic/Compositional Conversions (cont)

Tangent Hyperbolic	$\operatorname{arctanh}(x) = \operatorname{arctanh}(\sinh(x/\sqrt{(1+x^2)}))$
--------------------	---

Hyperbolic-Circular	$\operatorname{arctanh}(x) = \operatorname{arctanh}(\sinh(x/\sqrt{(1+x^2)}))$
---------------------	---

Circular 1	$\operatorname{arctanh}(x) = \operatorname{arctanh}(\sinh(x/\sqrt{(1+x^2)}))$
------------	---

Sine Cofunctional 1	$\operatorname{arcsinh}(x) = \operatorname{arctanh}(x/\sqrt{(1+x^2}))$
---------------------	--

Sine Cofunctional 2	$\operatorname{arcsinh}(x) = \pm \operatorname{arccosh}(\sqrt{1+x^2})$
---------------------	--


Cosine Cofunctional 1	$\operatorname{arccosh}(x) = \operatorname{arcsinh}(\sqrt{(x^2-1)}) , x \geq 1$
-----------------------	--

Cosine Cofunctional 2	$\operatorname{arccosh}(x) = \operatorname{arctanh}(\sqrt{(x^2-1)}/x) , x \geq 1$
-----------------------	--

Tangent Cofunctional 1	$\operatorname{arctanh}(x) = \operatorname{arcsinh}(x/\sqrt{(1-x^2)})$
------------------------	--

Tangent Cofunctional 2	$\operatorname{arctanh}(x) = \operatorname{arccosh}(1/\sqrt{(1-x^2)})$
------------------------	--

Unit Hyperbola



$$x^2 - y^2 = 1$$



By CROSSANT (CROSSANT)
cheatography.com/crossant/

Published 26th August, 2024.
Last updated 30th May, 2025.
Page 9 of 9.

Sponsored by [CrosswordCheats.com](https://crosswordcheats.com)
Learn to solve cryptic crosswords!
[http://crosswordcheats.com](https://crosswordcheats.com)