


### Introduction

Tool can be used to evaluate presence of neuropathic pain. Ask the older adult the questions below and score as noted. Higher scores are more indicative of pain with a neuropathic component. A score of 3 or higher indicates likely presence of neuropathic pain and justifies a more detailed evaluation. If the older adult has more than one painful area, they are to consider the one area that is most relevant to them. Conditions that might have a neuropathic pain component include diabetic or peripheral neuropathy, back pain, post-herpetic neuralgia, complex regional pain syndrome, leg/foot pain, large joint pain, and fibromyalgia.. Neuropathic Pain often requires different treatment approaches including use of anticonvulsants, antidepressants and other adjuvant medications.

Reproduced from Portenoy R for the ID Pain Steering Committee. (2006). Development and testing of a neuropathic pain screening questionnaire: ID Pain. Current Medical Research and Opinions, 22(6), 1555-1565.

### Pain Diagram



### Score

Question	Yes/No
1. Did the pain feel like pins and needles?	<input type="checkbox"/>
2. Did the pain feel like hot/burning?	<input type="checkbox"/>
3. Did the pain feel like numb?	<input type="checkbox"/>
4. Did the pain feel like electrical shocks?	<input type="checkbox"/>
5. Is the pain made worse with the touch of clothing or bed sheets?	<input type="checkbox"/>
6. Is the pain limited to your joints?	<input type="checkbox"/>

C

By [deleted]  
[cheatography.com/deleted-2754/](https://cheatography.com/deleted-2754/)

Published 14th April, 2016.  
 Last updated 12th May, 2016.  
 Page 1 of 1.

Sponsored by **ApolloPad.com**  
 Everyone has a novel in them. Finish Yours!  
<https://apollopad.com>