

### Introduction

NANDA International (NANDA-I), the Nursing Interventions Classification (NIC) and the Nursing Outcomes Classification (NOC) are comprehensive, research-based, standardized classifications of nursing diagnoses, nursing interventions and nursing-sensitive patient outcomes. The Nursing Diagnosis are used to develop the patient Care Plan, which contain the Interventions ((tasks) need to produce the expected Outcomes.

Credit:

### NIC codes 1-25

01. Activity intolerance
02. Ineffective Airway clearance
03. Latex Allergy response
04. Risk for latex Allergy response
05. Anxiety
06. Risk for Aspiration
07. Risk for impaired parent/infant/child Attachment
08. Disturbed Body image
09. Bowel incontinence
10. Ineffective Breastfeeding
11. Ineffective Breathing pattern
12. Decreased Cardiac output
13. Caregiver role strain
14. Impaired Comfort
15. Impaired verbal Communication
16. Parental role Conflict
17. Acute Confusion
18. Chronic Confusion
19. Constipation
20. Ineffective Coping
21. Compromised family Coping
22. Ineffective Denial
23. Diarrhea
24. Disturbed Energy field
25. Adult Failure to thrive

### NIC codes 26-50

26. Risk for Falls
27. Dysfunctional Family processes: alcoholism
28. Fatigue
29. Fear
30. Deficient Fluid volume
31. Excess Fluid volume
32. Impaired Gas exchange
33. Grieving
34. Anticipatory Grieving
35. Dysfunctional Grieving
36. Delayed Growth and development
37. Ineffective Health maintenance
38. Hopelessness
39. Hyperthermia
40. Functional urinary Incontinence
41. Total urinary Incontinence
42. Risk for Infection
43. Risk for Injury
44. Deficient Knowledge (specify)
45. Readiness for enhanced Knowledge (specify)
46. Impaired Memory
47. Impaired physical Mobility
48. Nausea
49. Imbalanced Nutrition: less than body requirements
50. Imbalanced Nutrition: more than body requirements

### NIC codes 51-75

51. Impaired Oral mucous membrane
52. Acute Pain
53. Chronic Pain
54. Impaired Parenting
55. Risk for Peripheral neurovascular dysfunction
56. Post-trauma syndrome
57. Powerlessness
58. Impaired Religiosity
59. Bathing/hygiene Self-care deficit
60. Feeding Self-care deficit
61. Risk for situational low Self-esteem
62. Disturbed Sensory perception specify: visual, auditory, kinesthetic, gustatory, tactile, olfactory

### NIC codes 51-75 (cont)

63. Impaired Skin integrity
64. Disturbed Sleep pattern
65. Spiritual distress
66. Risk for Suicide
67. Delayed Surgical recovery
68. Impaired Swallowing
69. Ineffective Therapeutic regimen management
70. Disturbed Thought processes
71. Impaired Tissue integrity
72. Ineffective Tissue perfusion specify type: renal, cerebral, cardiopulmonary, gastrointestinal, peripheral
73. Impaired Urinary elimination
74. Urinary retention
75. Wandering