

Background

- Not an inflammatory condition
- Occurs in synovial joints
- Due to genetic factors, overuse, and injury

Presentation

- Joint pain
- Stiffness for <30mins - worse with activity
- Crepitus
- Limited range of movement

Commonly Affected Joints

- Hips
- Knees
- SI joints
- DIPs + MCP
- Wrist
- Cervical spine


Complications of THR

- Neurovascular injury (external iliac)
- Infection
- Dislocation
- Periprosthetic fracture
- Loosening of implant + polyethylene wear, implant failure

Risk Factors


- Obesity
- Age
- Occupation
- Trauma
- Female
- Family history

Hand Signs




- Heberden's nodes (DIP)
- Bouchard's nodes (PIP)
- Squaring of thumb (CMC)
- Weak grip
- Reduced range of motion

Difference Between OA and RA



Xray Changes (LOSS)



Diagnosis

- No investigations needed if:
 - Patient over 45
 - Has typical activity-related pain
 - No morning stiffness
 - Stiffness lasts <30 mins

Management

- 1st line: Patient education, physio, OT, orthotics
- 2nd line: **Analgesia**
PO paracetamol + topical NSAIDs
Add PO NSAIDs (+ PPI)
Opiates (codeine, morphine)
- 3rd line: Intra-articular steroid injections
- 4th line: Joint replacement

C

By **ellieacook**
cheatography.com/ellieacook/

Published 29th July, 2022.
Last updated 28th July, 2022.
Page 1 of 1.

Sponsored by **Readable.com**
Measure your website readability!
<https://readable.com>