

### Definitions

**Electronegativity** is the ability of an atom to attract a pair of electrons towards itself in a covalent bond

**Ionic bonding** is the electrostatic attraction between oppositely charged ions (positively charged cations and negatively charged anions)

**Metallic bonding** is the electrostatic attraction between positive metal ions and delocalised electrons

**Covalent bonding** is electrostatic attraction between the nuclei of two atoms and a shared pair of electrons

**Bond energy** is the energy required to break one mole of a particular covalent bond in the gaseous state

**Bond length** is the internuclear distance of two covalently bonded atoms

**Sigma bond** is the covalent bond formed by 'head on' overlap of atomic orbitals


**Pi Bond** is the covalent bond formed by s- sideways overlap of atomic orbitals

**Dipole-Dipole** Forces are intermolecular attractions between molecules which are permanently polarised

**Hydrogen Bond** is the electrostatic attraction between a hydrogen atom which is bonded to a very electronegative atom and an electronegative atom of a neighbouring molecule

**Van Der Waals Forces** are intermolecular forces of attraction which arise from temporary dipoles in molecules

### Dative bonding



### Properties of water

**High melting & boiling points** Water has high melting and boiling points which is caused by the strong intermolecular forces of hydrogen bonding between the molecules

**High surface tension** Surface tension is the ability of a liquid surface to resist any external forces. The water molecules at the surface of liquid are bonded to other water molecules through hydrogen bonds. These molecules pull downwards on the surface molecules causing the surface to become compressed and more tightly together at the surface

### Properties of water (cont)

**Density** In ice the water molecules are packed in a 3D hydrogen-bonded network in a rigid lattice. Each oxygen atom is surrounded by hydrogen atoms. This way of packing the molecules in a solid and the relatively long bond lengths of the hydrogen bonds means that the water molecules are slightly further apart than in the liquid form

Hydrogen bonding in water, causes it to have anomalous properties such as high melting and boiling points, high surface tension and anomalous density of ice compared to water

### Factors influencing electronegativity

Nuclear charge	Increase in nuclear charge increases electronegativity
Atomic radius	Increase in atomic radius decreases electronegativity
Shielding by inner shells	Shielding causes decrease in electronegativity

### Paulings electronegativity to predict bonds

Paulings electronegativity	Bond
< 1.0	Covalent
1.0 - 2.0	Polar covalent
> 2.0	Ionic



By **Emergencybuttonse**

[chemistrybuttonse.com/emergencybuttonse/](https://www.chemistrybuttonse.com/emergencybuttonse/)

Published 15th September, 2024.

Last updated 15th September, 2024.


Page 1 of 3.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>

### Metallic bonding



In a metal, atoms are packed together in a lattice

Electrostatic attraction increases with:

Increase in positive charge

Decrease in size of metal ions

Increase in number of mobile electrons

### $\sigma$ and $\pi$ bonds

A pi bond is weaker than a sigma bond because the overlapping of charge clouds is less than in a sigma bond

H<sub>2</sub> has 1  $\sigma$  bond

C<sub>2</sub>H<sub>6</sub> has only  $\sigma$  bonds

C<sub>2</sub>H<sub>4</sub> has 1  $\sigma$  and 1  $\pi$  bond

HCN has 1  $\sigma$  and 2  $\pi$  bonds

N<sub>2</sub> has 1  $\sigma$  and 2  $\pi$  bonds

### VSEPR Theory and Molecular Shapes

Number of electron pairs	Electron pair geometries: 0 lone pair	1 lone pair	2 lone pairs	3 lone pairs	4 lone pairs
2	Linear 180°				
3	Trigonal planar 120°	Bent or angular <120°			
4	Tetrahedral 109°	Trigonal pyramidal <109°	Bent or angular <<109°		
5	Trigonal bipyramidal 120°/90°	Sawhorse or seesaw <120°/90°	T-shape 90°	Linear 180°	
6	Octahedral 90°	Square pyramidal <90°	Square planar 90°	T-shape 180°/90°	Linear 180°

### Van der Waals' Forces & Dipoles


Id - id forces increase with

Increasing number of electrons (and atomic number) in the molecule

Increasing the places where the molecules come close together

For small molecules with the same number of electrons, pd - pd forces are stronger than id - id

### Order of bond strength



### Hybridisation: sp, sp<sup>2</sup>, sp<sup>3</sup>

sp two different bonds (one may be triple bond)

sp<sup>2</sup> three different bonds (one may be double bond)

sp<sup>3</sup> four different bonds

### Hydrogen bonding

Hydrogen bonding is the strongest form of intermolecular bonding

For hydrogen bonding to take place the following is needed:

A species which has an O or N (very electronegative) atom with an available lone pair of electrons

A species with an -OH or -NH group

For hydrogen bonding to take place, the angle between the -OH/-NH and the hydrogen bond is 180°

### Bond Energy and Bond Length: Reactivity

Shorter bond length generally means higher bond energy, making molecules less reactive

Triple bonds are the shortest and strongest covalent bonds due to the large electron density between the nuclei of the two atoms

Triple bonds are the shortest and strongest covalent bonds due to the large electron density between the nuclei of the two atoms


The reactivity of a covalent bond is greatly influenced by: The bond polarity, the bond strength, the bond type ( $\sigma/\pi$ )

### Electronegativity relations

	Down a Group	Across a Period
Nuclear charge	Increases	Increases
Shielding	Increases	Reasonably constant
Atomic radius	Increases	Decreases
Electronegativity	Decreases	Increases


### Ionic bonding

Formation of sodium chloride (NaCl)




Ionic bonds are non-directional, each cation will attract any neighbouring anion and vice versa to form a huge ionic lattice

### Covalent bonding dot and cross




### Special octets

In expanded octet species, the central atom has **more** than eight electrons. An example is phosphorus(V) chloride, PCl<sub>5</sub>. This is possible only for Period 3 elements and beyond, this is because starting from Period 3, the atoms have **empty d orbitals** in the third energy level to accommodate more than eight electrons.



Octet deficient



Expanded octet

If the central atom is from Period 2 of the Periodic Table, the total number of electrons surrounding it cannot exceed eight (but can be less than eight).

If the central atom is from Period 3 and beyond, the total number of electrons surrounding it can exceed eight.



By **Emergybuttonse**

[cheatography.com/emergybuttonse/](https://cheatography.com/emergybuttonse/)

Published 15th September, 2024.

Last updated 15th September, 2024.

Page 2 of 3.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>