

Functions (cont)		Functions (cont)		Proofs (cont)		Proofs (cont)	
Inverse	has to be bijective. $f^{-1}(y) = x$ if and only if $f(x) = y$. because this is both 1to1 and onto, its a bijection, therefore invertible.	ex) let $f(0) = 0, f(1) = 0. f(2) = 2, f(3) = 4$	$f(x) = \text{floor}((x^2 - 2)/2)$. find $f(S)$ if $S = \{0, 1, 2, 3\}$	Proof by contradiction	Assume $\sim p$ is true, find contradiction, therefore $\sim p$ is true. prove that p is true if we can show that $\neg p \rightarrow (r \wedge \neg r)$ is true for some proposition r	UNIQUENESS proof	When asked for unique, prove exists, then unique. ex: $x \neq y$, so y doesnt have that property, therefore x is unique.
composition of fns	$f(g(a))$ or $f \circ g(a)$	equal functions	Two functions are equal when they have the same domain, the same codomain and map each element of the domain to the same element of the codomain	Counterexample	to show that a statement of the form $\forall x P(x)$ is false, we need only find a counterexample.	without loss of generality	an assumption in a proof that makes it possible to prove a theorem by reducing the number of cases to consider in the proof
floor/ceiling	bracketwithlow only/highonly. round down/up to nearest integer. ex) $-2.2 \text{ floor} = -3. 5.5 \text{ ceil} = 6$.			Proof by exhaustion	ex: Prove that $(n + 1)^3 \geq 3n$ if n is a positive integer with $n \leq 4$. Prove by doing $n = 1, 2, 3, 4$		
properties	$x \text{ floor} = n$ if and only if $n \leq x < n + 1$. $x \text{ ceil} = n$ if and only if $n - 1 < x \leq n$. $x \text{ floor} = n$ if and only if $x - 1 < n \leq x$. $x \text{ ceil} = n$ if and only if $x \leq n < x + 1$. $x - 1 < \text{floor}x \leq x \leq \text{ceil}x < x + 1$. $\text{floor}x = -\text{ceil}x$. $\text{ceil}x = -\text{floor}x$. $\text{ceil}(x + n) = \text{ceil}x + n$. opp of last floor						
		Proofs					
		Direct Proof	assume p is true, prove q . $p \Rightarrow q$. Always start with this then try contraposition.	Proof by cases	ex: Prove that if n is an integer, then $n^2 \geq n$. Case (i): When $n = 0$. Case (ii): When $n \geq 1$. Case (iii): In this case $n \leq -1$		
		Proof by contradiction	assume $\sim q$ is true, prove $\sim p$. ($\sim q \Rightarrow \sim p$) equals $(p \Rightarrow q)$	Constructive Existence Proof	$\exists x P(x)$. To find if $P(x)$ exists, show an example $P(c) = \text{True}$		
		Vacuous proof	if we can show that p is false, then we have a proof, called a vacuous proof, of the conditional statement $p \rightarrow q$	Nonconstructive Existence Proof	Assume no values makes $P(x)$ true. Then contradict.		
						Sets	
						Element of set	$a \in A, a \notin A$
						roster method	$V = \{a, e, i, o, u\}, O = \{1, 3, 5, 7, 9\}$.
						set builder notation	ex: the set O of all odd positive integers less than 10 can be written as: $O = \{x \in \mathbb{Z}^+ \mid x \text{ is odd and } x < 10\}$. ex) $A = \{x \mid x \geq -1 \wedge x < 1\}$. ex) $\{x \mid P(x)\}$
						Interval Notation	$[-2, 8)$
						Natural numbers	$\mathbb{N} = \{0, 1, 2, 3, \dots\}$
						\mathbb{N}	



Sets (cont)	
Integers Z	$Z = \{ \dots, -2, -1, 0, 1, 2, \dots \}$
Positive Integers Z^+	$Z^+ = \{1, 2, 3, \dots\}$
Rational Numbers Q	$Q = \{p/q \mid p \in Z, q \in Z, \text{ and } q \neq 0\}$
Real Numbers R	All previous sets (N, Z, Q)
R^+	positive real numbers
Complex numbers C	$\{a+bi, \dots\}$
Equal Sets	Two sets are equal if and only if they have the same elements. Therefore, if A and B are sets, then A and B are equal if and only if $\forall x(x \in A \leftrightarrow x \in B)$. We write $A = B$ if A and B are equal sets. Dont matter if its $\{1,3,3,3,2,2,3,\}$, still $\{1,3,2\}$. Also dont matter order.
Null/Empty Set	\emptyset , nothing. $\{\}$.
$\{\emptyset\}$	1 element
Singleton set	One element.

Sets (cont)	
Universal Set U	Universe in context of statement. Example vowels in alphabet: $U = \{z,y,x,w, \dots\}$, $A = \{a,e,i,o,u\}$ A is a subset of U.
Subset	$\forall x(x \in A \rightarrow x \in B)$. Ex) the set A is a subset of B if and only if every element of A is also an element of B. We use the notation $A \subseteq B$ to indicate that A is a subset of the set B. Ex) $A = \{1,2,3\}$, $B = \{1,2,3,4\}$, $A \subseteq B$.
Showing that A is a Subset of B	To show that $A \subseteq B$, show that if x belongs to A then x also belongs to B.
Showing that A is Not a Subset of B	To show that $A \not\subseteq B$, find a single $x \in A$ such that $x \notin B$.
Showing Two Sets are Equal	To see if $A = B$, Show $A \subseteq B$ and $B \subseteq A$

Sets (cont)	
proper subset	$\forall x(x \in A \rightarrow x \in B) \wedge \exists x(x \in B \wedge x \notin A)$ $A \subset B$ but $A \neq B$. B contains an element not in A. Ex) $A = \{1,2,3\}$, $B = \{1,2,3,4\}$. 4 makes it proper subset.
Cardinality	$ A $ Distinct elements of set. $A = \{1,2,3,3,4,4\}$ $ A = 4$
Power Set	the power set of S is the set of all subsets of the set S. The power set of S is denoted by $P(S)$. Ex) $A = \{1,2,3\}$. $P(A) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\}$. Ex) $P(\{\emptyset\}) = \{\emptyset, \{\emptyset\}\}$
Cardinality of Power Set	2^n , n is elements.
Tuple	$(a_1, a_2, a_3, \dots, a_n)$ Ordered. Ex) $(5,2) \neq (2,5)$

Sets (cont)	
Cartesian Product	$\{(a, b) \mid a \in A \wedge b \in B\}$. The Cartesian product of A and B, denoted by $A \times B$, is the set of all ordered pairs (a, b), where $a \in A$ and $b \in B$. Ex) $A = \{0,1\}$ $B = \{2,3,4\}$, $A \times B = \{(0,2),(0,3),(0,4), (1,2),(1,3),(1,4)\}$
Truth Set	$P(x): \text{abs}(x) = 3$. Truth Set of $P(x) = \{3,-3\}$

Set Operations	
Union	$A \cup B = \{x \mid x \in A \vee x \in B\}$. Ex) $A = \{1,4,7\}$ $B = \{4,5,6\}$. $A \cup B = \{1,4,5,6,7\}$
Intersection	$A \cap B = \{x \mid x \in A \wedge x \in B\}$. Ex) $A = \{1,4,7\}$ $B = \{4,5,6\}$. $A \cap B = \{4\}$.
disjoint	If $A \cap B = \text{nothing}$, A and B are disjoint.



Set Operations (cont)

principle of inclusion-exclusion
 $|A \cup B| = |A| + |B| - |A \cap B|$
 ex) $A = \{1,2,3,4,5\}$, $B = \{4,5,6,7,8\}$.
 $|A \cup B| = 5 + 5 - 2 = 8$

$A - B$, difference of A and B
 $A - B = A \cap B^c = \{x \mid x \in A \wedge x \notin B\}$
 Elements in A that are not in B.
 Ex) $\{1, 3, 5\} - \{1, 2, 3\} = \{5\}$. This is different from the difference of $\{1, 2, 3\}$ and $\{1, 3, 5\}$, which is the set $\{2\}$.

Complement of A, A^c
 $\{x \in U \mid x \notin A\}$
 Everything in the universe context that's not in A.
 Ex) $U = \{1,2,3,4\}$.
 $A = \{2\}$ $B = \{3\}$.
 $A^c = \{1,3,4\}$

$U = \mathbb{R}$
 $A = \{x \mid x \geq -1 \wedge x < 1\}$
 $B = \{x \mid x < 0 \vee x \geq 2\}$
 $A \cup B = \{x \mid \square < 1 \vee x \geq 2\}$.
 $A \cap B = \{x \mid x < 0 \wedge \square \geq -1\}$.
 $A^c = \{x \mid x < -1 \vee x \geq 1\}$.

Set Operations (cont)

Identity, , , , , , absorption,
 $A \cap U = A$. $A \cup \emptyset = A$.

domination
 $A \cup U = U$. $A \cap \emptyset = \emptyset$

idempotent
 $A \cup A = A$. $A \cap A = A$

complementation
 $(A^c)^c = A$

commutative
 $A \cup B = B \cup A$. $A \cap B = B \cap A$

associative
 $A \cup (B \cap C) = (A \cup B) \cap C$.
 $A \cap (B \cup C) = (A \cap B) \cup C$

distributive
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$.
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

de morgan's
 $(A \cap B)^c = A^c \cup B^c$.
 $(A \cup B)^c = A^c \cap B^c$

absorption
 $A \cup (A \cap B) = A$.
 $A \cap (A \cup B) = A$.

complement
 $A \cup A^c = U$. $A \cap A^c = \emptyset$.

Set Operations (cont)

countable
 a set that either is finite or can be placed in one-to-one correspondence with the set of positive integers. To be countable, there must exist a 1-1 and onto (bijection) between the set and $\mathbb{N}!$ (i.e. \mathbb{Z}^+)

Ex) Show that the set of positive even integers E is countable set.
 Let $f(x) = 2x$. Then f is a bijection from \mathbb{N} to E since f is both one-to-one and onto. To show that it is one-to-one, suppose that $f(n) = f(m)$. Then $2n = 2m$, and so $n = m$. To see that it is onto, suppose that t is an even positive integer. Then $t = 2k$ for some positive integer k and $f(k) = t$

