

by Jerstellar via cheatography.com/204102/cs/43565/

Module 1 - Characteristics of Life		
Characteristic	Example	
Growth and Development	cell growth and cell division	
Maintain Homeostasis	maintain appropriate concentrations of different chemicals, pH level, optimum temperature	
Reproduction	succeeding generations thruough sexual or asexual processes	
Response to Environment or Stimuli	adaptation to environment	
Energy Processing	photosynthetic process	
Organized	highly organized and coordinated cell structures	

Terms	
Specializ- ation	adaptation of an organ or part to serve a special function
Merist- hematic Tissue	includes undifferentiated cells that are capable of specialization; most action takes place here
Cytoki- nesis	physical separation of the two daughter cells; where cell division ends
Hibern- ation	state of minimal activity and metabolic depression
Torpor	involuntary and lasts for just a few hours during the daytime; not as heavy as hibernation
Piloer- ection	or goosebumps ; modification of the heat exchange, contraction of the musculi arrectores pilorumor (MAP)
Autotrophs	organisms that can make their own food
Photosynt- hetic process	+ Oxygenic photosynthesis - Light energy transfers electrons from water to carbon dioxide in order to produce carbohydrates; seen in algae, cynobacteria, plants

Terms (cont)	
	+ Anoxygenic photosynthesis - light energy is captured and converted to ATP, without the production of oxygen; doesn't have water as electron donor
Epithelial tissues	line the cavities and surfaces of the body such as the inside of the stomach and the outermost skin layer
Connective tissue	supports, protects and binds certain parts of the body such as muscles, together
Muscular tissues	produce movement by contraction and expansion
Nervous tissue	receive stimuli and conduct electrical impulses
Dermal tissue	forms outer covering of plants
Vascular tissue	moves water and nutrients through the plant
Ground tissue	makes up most of plants' bodies and performs majority of bodily functions
Vegetative organs	help sustain plant life; roots and leaves
Reprod- uctive organs	facilitate either sexual or asexual reproduction; cones, flowers and fruits
Shoot system	parts above the ground; leaves, and stems
Root system	parts below the ground; roots and tubers
Diurnality	behavior characterized by activity during daytime, highly variable temperature
Crepuscul- arity	active primarily during the twilight period, coldest temperature is at dawn



By **Jerstellar** cheatography.com/jerstellar/

Not published yet. Last updated 17th July, 2024. Page 1 of 4.



by Jerstellar via cheatography.com/204102/cs/43565/

Terms (cont)

Noctu behavior in animals characterized by being active during rnality the night and sleeping during the day, constant temperature

Human body has 11 systems, while plants only have 2 (shoot and root)

Life Processes		
Movement	transfer places with the use of specialized structures like flagella, cilia and pseudopodia	
Respir- ation	exhibits a metabolic pathway that breaks down glucose and produces adenosine triphosphate (ATP)	
Sensitivity	act on sensing a stimulus and at the same time responding to it	
Growth	increase the size of each individual cell or increase of the number of cells	
Reprod- uction	formation of new cells by the process of cell division to replace or repair old cells	
Excretion	get rid of by-products due to metabolic processes which maybe toxic	

break down food, into simple molecules that can be

absorbed and be utilized

Module 2 - Cell Theory

Nutrition

Development of	of Microscope
Zachrias Janssen	Dutch spectacle-maker who discovered the first compound microscope which was later disputed
Galileo Galilei	able to make his own microscope because of his knowledge about glass and focal lengths
Antony van Leeuwe- nhoek	"Father of Microbiology", able to discover bacteria and protozoa; called bacteria "animacules"
Robert Hooke	coined the term "cell"; published "Micrographia" in 1665
Formation of P	rostulates

Matthias Schleiden	German botanist, noticed that plants are made up of cells
Robert Schwann	concluded that animals are made up of cells; coined "- Schwann cells", which myelinates the axons of the peripheral nervous system
Rudolf Virchow	"Father of Modern Pathology", published "Virchow's archives" and the aphorism "every cell stems from another cell"

Postulates of Cell Theory

- 1. All known living things are made up of cells.
- 2. The cell is a structural and functional unit of all living things.
- 3. All cells come from pre-existing cells by division.

Additional:

1. All cells contain hereditary information which is passed from cell to cell

during division.

- 2. All cells are basically the same in chemical composition.
- 3. All energy flow of life occurs within cells.

Module 3 - Prokaryotic vs. Eukaryotic cells

All known life are classified into 3 domains: Archea, Bacteria, Eukarya. The organisms in **Archea and Bacteria are prokaryotes** while the organisms in **Eukarya have eukaryotic cells**.

Prokaryotic cells - simpler and lack the membrane-bound organelles and nucleus; more primitive than eukaryotes, single-celled.

Eukaryotic cells - single or multicelllar

Major Difference in Cell Structure

Eukaryotes store their DNA as chromosomes within the nucleus but prokaryotes lack the nucleus. Instead, the majority of their DNA is in the **nucleoid**. Additional DNA pieces, called **plasmids**, are shaped like rings and reside outside the nucleoid in the cytoplasm.

Differences in Organization

Eukaryotic cells use a specific cell division process called **mitosis**, while prokaryotic cells use **binary fission**.

- -Prokaryotes create an exact copy of themselves; though genetic variance occur through **transduction**, which is when virus transmit plasmid containing DNA to bacterial cells (host).
- -Eukaryotes sexually reproduce through **meiosis**, which maximizes genetic diversity and minimizes mutation.



By **Jerstellar** cheatography.com/jerstellar/

Not published yet. Last updated 17th July, 2024. Page 2 of 4.




by Jerstellar via cheatography.com/204102/cs/43565/

Other Differences

Features	Prokaryotic	Eukaryotic
Protective layers	Capsule, cell wall and cell	Cell membrane (animal
	membrane	cells), cell wall and cell
		membrane (plant cells)
	Usually unicellular (some	Usually multicellular
Cell type	cyanobacteria maybe	
	multicellular)	
Complexity	Simple	Complex organization
Nucleus location	Free in the cytoplasm,	Contained in membrane
Nucleus location	attached to mesosomes	bound structure
	Usually single circular	Multiple linear with
Chromosome	without histones	histones
	Expressed in groups	Expressed individually
Genes	called operons	
Genome	DNA haploid genome	DNA diploid genome
	Efficient and compact	With large amounts of
Genome nature	with little repetitive DNA	non-coding repetitive
		DNA
Movement	Simple flagellum, if	Complex flagellum, if
Movement	present	present
	Via cytoplasmic	Via mitochondria
Respiration	membrane	
	Electron transport chain	Within membrane bound
Energy production site	located in the cell	mitochondria
	membrane	
	Wide variation	Glycolysis, electron
Metabolic mechanism		transport chain, Krebs
		cycle
DNA replication	Occurs in cytoplasm	Occurs in the nucleus
	•	
	Occurs simultaneously	Transcription occurs in
Transcription and		nucleus and then
translation		translation occurs in
		cytoplasm

Similarities between Prokaryotes and Eukaryotes



Both have DNA, plasma membrane, ribosomes for protein synthesis, and cytoplasm

Terms	
Proteins	comprise hundreds or thousands of smaller units called amino acids (20 types)
Carboh- ydrates	provide energy, structural support and cellular communication; plant and fungal cell walls have carbohydrate cell walls
Lipids	made up of fatty acids that can either be saturated or unsaturated
Nucleic Acids	DNA and RNA
Archea	single-celled microorganisms living in environments low in oxygen (extremophiles)
Flagellum	specialized part used for movement
Cytoplasm	jelly-like fluid within in a cell that is composed primarily of water, salts and proteins
Ribosome	organelle used to synthesize proteins
Bacteria	organelle used to synthesize proteins

Terms (cont)

Operon a functioning unit of DNA containing a cluster of genes under the control of a single promoter

Module 4 - M	lembrane-bound Organelles
Nucleus	consists of nuclear envelope, chromatin and nucleolus; largest and contains genome
Endopl- asmic Reticulum	major site of synthesis, flattened sac network (cisternae). Its function is closely linked to that of the golgi apparatus and together they form the cell's secretory route
Rough Endopl- asmic Reticulum	takes proteins from the cytosol and continues its production in the golgi apparatus until completion
Smooth Endopl- asmic Reticulum	lipid, phospholipid and steroid synthesis
Golgi Apparatus	packages macromolecules into vesicles; modifies proteins and lipids from endoplasmic reticulum
Mitoch- ondria	site of ATP synthesis; helps maintain the intracellular environment, has inner and outer membrane with an intermembrane space in between
Lysosomes	acidic; contain numerous hydrolytic enzymes which catalyze hydrolysis reactions
Perixo- somes	contain the enzyme catalase which decomposes hydrogen peroxide
Vacuoles	acts as a storage for nutrients as well as waste materials to protect the cell from toxicity; helps in maintaining an acidic internal pH



By **Jerstellar** cheatography.com/jerstellar/

Not published yet. Last updated 17th July, 2024. Page 3 of 4.



by Jerstellar via cheatography.com/204102/cs/43565/

Module 4 - Membrane-bound Organelles (cont)		
Vesicles	facilitate the storage and transport of materials in and outside the cell	
Chloro- plast	produces amino acids and lipids required for the production of chloroplast membrane; has two distinct regions- grana and stroma	

Terms	
Cell compartme- ntalization	process of selectively permeable nuclear envelope (separates the contents of the nucleus from the cytoplasm)
Gene expression	involves first transcription, which is the mechanism by which DNA is transcribed to mRNA
pre-mRNA	undergoes a process known as post-transcriptional modification where molecules are added or removed
Cytochrome p450	enzyme in SER; essential to some drugs and toxins, such as alcohol and barbiturates, in the metabolism
Exocytosis	form of active transport and bulk transport in which a cell transports molecules out of the cell
Protein processing	carbohydrate regions of glycoproteins are altered by addition, removal or modification of carbohydrates
Lipid processing	adds phosphate groups and glycoproteins to lipids from ER (like cholesterol) to create the phospholipids that make up the cell membrane

Terms (cont)	
Mannose 6-phos- phate receptor	lysosomal protein; binds newly synthesized lysosomal hydrolases in the trans-Golgi network and deliver them to pre-lysosomal compartments
Secretory proteins	hormones; requires ATP, as it is necessary to fuse two negatively charged membranes to allow its release
Cell-s- urface proteins	phospholipids; primary route of communication among the cells and the external environment
Porins	protein in the outer membrane of nucleus; enable ion movement to and from mitochondrion
Phagoc- ytosis	important in killing mechanisms which are independent of oxygen
Tonoplast	membrane which bounds the vacuole of a plant cell
Lamellar phase	similar to plasma membrane; outer layer enclosing the liquid in vesicle
Grana	made up of thylakoids; sight for the process of light dependent reactions of the photosynthesis process
Stroma	contains grana and is similar to the cytoplasm in cells in which all the organelles are embedded
Stroma Lamellae	flat membranous tubules; connects the thylakoids of the different grana

Module 5 - Non-Membrane-bound Organelles

Ribosomes



By **Jerstellar** cheatography.com/jerstellar/

Not published yet. Last updated 17th July, 2024. Page 4 of 4.