# Cheatography

## NumPy by Justin1209 (Justin1209) via cheatography.com/101982/cs/21247/

#### Import Statement

import numpy as np

#### **Creating Arrays**

```
# Create a numpy array
array 1 = np.array([92, 94, 88,
91, 87])
# Create a numpy array from a
CSV
test_2 = np.genfromtxt('test -
2.c sv', delimi ter =',')
# Create a two-di men sional
array
test_1 = np.arr ay([92, 94, 88,
91, 87])
test 2 = np.arr ay([79, 100, 86,
93, 91])
test 3 = np.arr ay([87, 85, 72,
90, 92])
np.arr ay( [[92, 94, 88, 91,
87],
                   [79, 100, 86,
93, 91],
                   [87, 85, 72,
90, 92]])
```

#### Operations with Arrays

```
arr = [1, 2, 3, 4, 5]
# Adding 3 to each entry
>>> a = np.arr ay(arr)
>>> a plus 3 = a + 3
# Adding arrays
>>> a = np.arr ay([1, 2, 3, 4,
51)
>>> b = np.arr ay([6, 7, 8, 9,
10])
>>> c = a + b
# Logical Operations
>>> a = np.arr ay([10, 2, 2, 4,
5, 3, 9, 8, 9, 7])
>>> a > 5
```

By Justin1209 (Justin1209)

cheatography.com/justin1209/

```
Operations with Arrays (cont)
```

```
> array([True, False, False, False, False,
False, True, True, True, True], dtype=bool)
>>> a[a > 5]
array([10, 9, 8, 9, 7])
>>> a[(a > 5) | (a < 2)]
array([10, 9, 8, 9, 7])
-> c: array([ 7, 9, 11, 13, 15])
```

### Selecting from Arrays (1 Dimension)

```
a = np.array([5, 2, 7, 0, 11])
>>> a[0]
-> 5
>>> a[-1]
-> 11
>>> a[-2]
-> 0
>>> a[0:5:2]
-> *array([5, 7, 11])
>>> a[1:3]
-> array([2, 7])
>>> a[:3]
-> array([5, 2, 7])
>>> a[-3:]
-> array([7, 0, 11])
```

Selecting from Arrays (2 Dimensions)

```
-> Basic Procedure a[row, column]
a = np.arr ay([32, 15, 6, 9,
14],
                            [12,
10, 5, 23, 1],
                            [2,
16, 13, 40, 37]])
# selects the first column
>>> a[:,0]
-> array([32, 12, 2])
# selects the second row
>>> a[1,:]
-> array([12, 10, 5, 23, 1])
# selects the first three
elements of the first row
>>> a[0,0:3]
-> array([32, 15, 6])
```

### Selecting Elements

```
np.count_nonzero(poodle_colors
== "brown")
-> returns the number of poodles
with brown hair
```

Mean and Logical Operations (On arrays)

```
np.mean(array > 8)
-> returns the percentage of
values in the array that meet
the criteria
```

We can use np.mean to calculate the percent of array elements that have a certain property.

Published 28th November, 2019. Last updated 18th December, 2019. Page 1 of 2.

Sponsored by Readable.com Measure your website readability! https://readable.com

### NumPy

### by Justin1209 (Justin1209) via cheatography.com/101982/cs/21247/

Mean over 2 Dimensional Arrays
>>> ring_toss = np.array([[1, 0,
0],
-
[0, 0, 1],
-
[1, 0, 1]])
>>> np. <b>mean</b> (ring_ toss)
0.44 -> Overall Average
>>> np.mean(ring_ toss,axis=1)
array([ 0.33, 0.33, 0.67]) ->
Average per row
>>> np.mean(ring_ toss,axis=0)
array([ 0.67, 0. , 0.67]) ->
Average per column

Cheatography

### Dealing with Outliers

# Sort the Dataset
np.sort(array)
-> Outliers are clearly visible
now

#### Percentiles

d = np.array([1, 2, 3, 4, 4, 4, 6, 6, 7, 8, 8]) np.percentile(d, 40) -> 4.00

#### Shape (dimensions) of an array

The **.shape** attribute for NumPy arrays returns the dimensions of the array. If array has n rows × m columns, then array.shape returns (n, m). # Generate own Normal

Distribution Set
-> np.ran dom.no rma l(loc,
scale, size)
loc: the mean for the normal
distri bution
scale: the standard deviation of
the distri bution
size: the number of random
numbers to generate

68% of our samples will fall between +/- 1 standard deviation of the mean

95% of our samples will fall between +/- 2 standard deviations of the mean

99.7% of our samples will fall between +/- 3 standard deviations of the mean

### **Binomial Distribution**

```
np.random.binomial (N, P, size)
N: The number of samples or
trials
P: The probab ility of success
size: The number of experi ments
#Basketball Example
  Let's generate 10,000 " exp -
eri men ts"
N = 10 shots
P = 0.30 (30% he'll get a free
throw)
-> a = np. random.bi nomia¥10,
0.3, 10000)
# Probab ility that he makes 4
Shots:
prob = np.mean(a == 4)
```

The **binomial distribution** can help us. It tells us **how likely** it is for a **certain number of "successes"** to happen, given a probability of success and a number of trials.



By Justin1209 (Justin1209) cheatography.com/justin1209/

Published 28th November, 2019. Last updated 18th December, 2019. Page 2 of 2. Sponsored by Readable.com Measure your website readability! https://readable.com