

Libraries to Import

<code>library(tidyverse)</code>	<code>library(broom)</code>
<code>library(mosaic)</code>	<code>library(dplyr)</code>

Block 2 Quiz 1

mutually exclusive	two cannot be concurrent
independent	no relevance/influence between 2 factors
complement	opposite
<code>rbinom(# of trials, # of attempts within trial, success rate) mean(results > amt)</code>	returns probability of success that the amt is exceeded
<code>mean(rbinom(simulatedTrials, #ofAttempts, successRate))</code>	
<code>favstats(results)</code>	finding 3rd quartile of probability

`quantile(results, 0.75)`

`qbinom(0.75, size = #ofAttempts, prob = successRate)`

`choose(n, k)` n choose k. different ways for result

Block 1 GR

Categorical	non-number
Numerical	can be categorized by number
continuous	non-whole #
discrete	whole #
stratified sampling	some of all groups

Block 1 GR (cont)

cluster sampling one group, all variables

simple random sampling random selection across all groups

observational study not experimenting

retrospective study data collected after the fact

prospective study data collected during event

skew tail is what skew it is

`favstats(colforeach~thiscol, data = dataSet)` returns min, max, SD, mean, median

explanatory variable x

response variable y

independence has no trend

`tally(~col1+col2, data = dataSet, format = "percent/decimal")` % of the dataset that applies to both

`tally((colforeach~thiscol, data = dataSet, format = "percent", margin = TRUE))` of col1, how many also in col2?

`dataSet%>% gf_props(~Survived, fill = ~Pclass, position = "fill")%>% gf_labs(title = "", subtitle = "", x = "", y = "")%>% gf_theme(theme_bw())`

Block 1 GR

Categorical non-number

Numerical can be categorized by number

continuous non-whole #

discrete whole #

stratified sampling some of all groups

cluster sampling one group, all variables

simple random sampling random selection across all groups

observational study not experimenting

retrospective study data collected after the fact

prospective study data collected during event

skew tail is what skew it is

`favstats(colforeach~thiscol, data = dataSet)` returns min, max, SD, mean, median

explanatory variable x

response variable y

independence has no trend

`tally(~col1+col2, data = dataSet, format = "percent/decimal")` % of the dataset that applies to both

`tally((colforeach~thiscol, data = dataSet, format = "percent", margin = TRUE))` of col1, how many also in col2?

`dataSet%>%`



By **kaylahs**

cheatography.com/kaylahs/

Not published yet.

Last updated 12th December, 2022.

Page 1 of 2.

Sponsored by **CrosswordCheats.com**

Learn to solve cryptic crosswords!

<http://crosswordcheats.com>

Block 2 Quiz 2

`pnorm(value, mean mu, stdev)` returns CDF of normal distribution

Probability of randomly selecting within a range

upper bound `pnorm`
- lower bound `pnorm`

`qnorm(percentile value, mu, stdev)` returns percentile of dataset

`integrate(function(x)- {functionInfo}, lowerBound, upperBound)$value`

Integrate

`adaptIntegrate(function, lowerLimit = c(lowerBound, lowerBoundY), upperLimit = c(upperBoundX, upperBoundY))$integral`

integrate with 2 variables

$Cov(X,Y) = E(XY) - E(X)E(Y)$

covariance

$Cov(X,Y) = 0$

Independence

CDF

cumulative density function. probability that a random variable will take on a value \leq given value. integral of PDF

PDF

probability density function. probability that a random variable will take on a given value. derivative of CDF.

Block 2 Quiz 2 (cont)

`rexp(# of trials, mu) + 1 ///` exponential distribution

`mean(results < givenNumber)`

`pexp(xValueNumber, rate = mu)` exponential distribution

Block 3

`lm()` im not done with this!

bootstrapping



By **kaylahs**

cheatography.com/kaylahs/

Not published yet.

Last updated 12th December, 2022.

Page 2 of 2.

Sponsored by **CrosswordCheats.com**

Learn to solve cryptic crosswords!

<http://crosswordcheats.com>