

### Fever - classes of medication

Acetaminophen **Mechanism of action:** acts at hypothalamus to cause peripheral vasodilation, which enables sweating and allows body to rid excess heat  
No anti-inflammatory action  
**Primary use:** fever, mild to moderate pain, osteoarthritis

NSAID's Same mechanism of action as acetaminophen (for fever)  
Because of acetaminophen's safety record (few drug interactions and side effects), it is first-line for fever  
NSAID could be more appropriate if inflammation is also present (ibuprofen > ASA)  
ASA is contraindicated in children Reye's Syndrome (ASA + virus + fever in child)

### Adverse effects

NSAID nausea, dyspepsia, ulcer with long-term use, potential anti-platelet action, hypertension, increased risk of cardiac event with long-term use  
Take with food  
Caution in kidney disease, cardiovascular disease, GI conditions

Corticosteroids

### Adverse effects (cont)

Acetaminophen very rare liver toxicity (max dose of 4g/24-hours), avoid alcohol, interacts with warfarin (but doesn't ↑ bleeding on its own)

1st-Generation Antihistamines Significant sedation

### 2nd-Generation Antihistamines

Intranasal Corticosteroids nasal irritation, dryness and bleeding (epistaxis), bad taste, loss of smell

Decongestants *Phenylephrine, pseudoephedrine* oral – hypertension, anxiety, insomnia; intranasal – nasal irritation, rebound congestion, rarely systemic effects

Penicillin anaphylaxis, diarrhea, nausea, vomiting, pain at injection site, superinfections, some (minor) drug interactions

Cephalosporins *Cefotaxime* (3rd Gen.) hypersensitivity, rash, itching, anaphylaxis, diarrhea, vomiting, nausea, pain at injection site, some (minor) drug interactions  
Must be given IV or IM (not orally)

### Adverse effects (cont)

Tetracyclines diarrhea, yeast infections, nausea, vomiting, epi-gastric burning, yellow-brown teeth discoloration in young children (we don't prescribe for kids), photosensitivity  
Can potentially interfere with oral contraceptives (recommend backup method)  
Higher chance of superinfections because it is broad-spectrum

Macrolides *Erythromycin* : significant nausea, vomiting, diarrhea (take with food), some important drug interactions  
Warfarin, cyclosporine, anticonvulsants (all via CYP450 inhibition/induction)  
Fidaxomicin – new; for treatment of *c. difficile*; not absorbed, stays in GI tract  
nausea, constipation, vomiting

Aminoglycosides *Gentamicin* ototoxicity, nephrotoxicity

By [kjaniskevich](https://kjaniskevich)



[cheatography.com/kjaniskevich/](https://cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 1 of 8.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Adverse effects (cont)

Fluoroquinolones  
*Ciprofloxacin*

nausea, diarrhea (can take with food), photosensitivity

Separate from minerals like calcium, iron, magnesium, aluminum (including supplements and antacids) by 2h

Serious adverse effects associated with fluoroquinolones: Tendinitis or tendon rupture (1.3-5.6 in 10,000)

Cardiac arrhythmias (15-57 in 100,000)

CNS effects

seizures, tremors, altered mental state

Peripheral neuropathy

Sulfonamides  
*Sulfamethoxazole-Trimethoprim*  
(SMZ-TMP, *Septra*®, *Bactrim*®, -DS)

nausea, vomiting, skin rashes, photosensitivity, anemia, crystalluria

Drink lots of water to prevent crystalluria

Monitor for: painful urination, abdominal pain, blood in urine, fever

Carbapenems  
*ertapenem*, *imipenem*, *meropenem*

skin reactions, inflammation at injection site, diarrhea, nausea, vomiting

Clindamycin

High risk of superinfection (GI)

### Adverse effects (cont)

Nitrofurantoin

Changes urine to orange colour

Must take with food

Metronidazole

Disulfiram reaction – flushing, tachycardia, shortness of breath, severe nausea & vomiting, throbbing headache, visual disturbance, confusion, dizziness

Occurs ~ 5-10 minutes after intake, lasts 30 mins several hours

Vancomycin

Ototoxicity and nephrotoxicity

Linezolid

lactic acidosis, myelosuppression (↓WBC and platelets), peripheral and optic neuropathy, serotonin syndrome, diarrhea,

Major drug interaction with any serotonergic drug, may need to discontinue until course of treatment finished, also inhibits MAO

Rifampin (RMP)

Rashes, blood dyscrasias, GI disturbances, liver damage, nephrotoxicity

Secretions coloured a reddish-orange (sweat, urine, sputum, tears)

### Adverse effects (cont)

Amphotericin B

fever & chills during infusion, vomiting, headache, phlebitis, nephrotoxicity, hypokalemia, ototoxicity

Azole Antifungals  
*fluconazole*, *itraconazole*, *ketoconazole*, *miconazole*, *voriconazole*

Rare hepatotoxicity – avoid alcohol, watch for jaundice, monitor liver enzymes

Nystatin

Oral thrush – swish and swallow oral suspension four times daily (works topically) (needs Rx)



By [kjaniskevich](https://www.cheatography.com/kjaniskevich)

[cheatography.com/kjaniskevich/](https://www.cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 2 of 8.

Sponsored by [ApolloPad.com](https://www.apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://www.apollopad.com>

### Adverse effects (cont)

Classic	Increased risk of infections,
Immuno	Increased risk of cancers such
-suppr-	as lymphomas, cysts, and polyps
essant	Frequency increases with
	intensity and duration of
	treatment, Kidney impairment,
	hepatic impairment, Hypert-
	ension, hyperlipidemia,
	CNS: tremor, headache, skin
	prickling sensation,
	GI: nausea, vomiting, abdominal
	pain, diarrhea, gingival
	hyperplasia,
	MSK: Muscle cramps, myalgia
	Endocrine: Menstrual distur-
	bances, gynecomastia, Hypert-
	richosis (abnormal amount of
	hair growth over body), Fatigue

### Adverse effects (cont)

Chemot	<b>short term</b>
herapy	Nausea/vomiting, Diarrhea or
	constipation, Mucositis/stoma-
	titis, Myelosuppression, Hair
	growth alterations, Weight gain /
	weight loss, Taste alterations,
	Fatigue, Hepatic and renal
	changes, Cardiac function
	changes, Rash / skin changes /
	nail changes, High blood
	pressure
	<b>Long term</b>
	Infertility, Secondary malign-
	ancies, Heart failure, Osteop-
	orosis, Pulmonary fibrosis,
	Cataracts, Peripheral neurop-
	athy, Hearing loss, Fatigue,
	Endocrine abnormalities

### Inflammation mediators

Histamines

Bradykinin

Leukotrienes

Cytokines

Interleukins

Prostaglandin

### Inflammation classes of medication

<b>Non-ste-</b>	Inhibit cyclo-oxygenase (COX),
<b>roidal</b>	which reduces prostaglandin
<b>anti-infl-</b>	synthesis therefore inhibiting
<b>ammato-</b>	inflammation
<b>ries</b>	Also have analgesic and
<b>NSAID</b>	antipyretic properties
	For mild to moderate inflam-
	mation

### Inflammation classes of medication (cont)

COX-1	– In all tissues, stomach
	lining (mucosa),
	involved in platelet
	aggregation
COX-2	more specific for inflam-
	mation
<i>Ibuprofen</i>	mild to moderate inflam-
	mation, fever, mild to
	moderate pain, dysmen-
	orrhoea, musculoskeletal
	pain, arthritis

<b>Corticosteroids</b>	Mimic endogenous
	cortisol, attempting to
	bring body back to
	homeostasis after a
	fight-or-flight response
	Anti-inflammatory and
	immuno-suppressive
	For severe inflammation
	Serious systemic
	adverse effects limit use
	to emergencies and
	severe inflammation
	(multiple sclerosis,
	rheumatoid arthritis,
	auto-immune diseases)
	Local administration,
	short-term use preferred
	whenever possible



By [kjaniskevich](https://www.cheatography.com/kjaniskevich/)

[cheatography.com/kjaniskevich/](https://www.cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 3 of 8.

Sponsored by [ApolloPad.com](https://www.apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://www.apollopad.com>

### Antibiotics - Classes of medication

<b>Penicillins</b> <i>-cillin</i>	Disrupt bacterial cell walls, Bactericidal Penicillin-binding protein: a protein only in bacterial cell walls that penicillin binds to which weakens the cell wall, allows fluid to enter and destroys the cell Penicillins contain a beta-lactam ring in its structure necessary for activity Many bacteria produce beta-lactamase (penicillinase) that is a natural defense to penicillin – it breaks the beta-lactam ring, leaving it ineffective therefore penicillin resistance
<i>Amoxicillin + Clavulanic Acid (Amoxiclav)</i>	Clavulanic acid inhibits $\beta$ -lactamases (penicillinases) of some microorganisms to allow amoxicillin to be active against it Synergistic relationship

### Antibiotics - Classes of medication (cont)

<i>Penicillin G (Pen G)</i>	Drug of choice against streptococci, pneumococci, staphylococci, gonorrhea and syphilis (given IV or IM)
<b>Cephalosporins</b> <i>-ce(f)ph</i>	Related to penicillins (1st gen. also have beta-lactam ring) Also inhibit cell wall synthesis, Bactericidal Classified according to “generation” (1 - 4) <b>General Rules</b> 1st generation not effective against bacteria producing beta-lactamase More potent as go up in generation Fewer similarities with penicillins as go up in generation Higher generations reserved for known resistant infections
<i>Cefotaxime (3rd Gen.)</i>	Has broad-spectrum activity against gram-negative organisms; for serious infections of lower respiratory tract, CNS, genitourinary system, bones, blood, and joints

### Antibiotics - Classes of medication (cont)

<b>Tetracyclines</b> <i>doxycycline, minocycline, tetracycline, -cycline</i>	Inhibit bacterial protein synthesis, Bacteriostatic Broad-spectrum (both gram-positive and negative) Usually given orally (PO) Should not be given at the same time as iron, calcium, magnesium (ions bind to drug so it can't absorb) – separate by 2h
<i>Tetracycline</i>	Used for Rocky Mountain spotted fever, h.pylori infections, acne vulgaris, chlamydia
<b>Macrolides</b> <i>azithromycin, clarithromycin, erythromycin, fidaxomicin, -thromycin</i>	Inhibit bacterial protein synthesis, Some are bactericidal, some bacteriostatic No structural similarities to penicillin – zero chance of cross-reactivity
<i>Erythromycin</i>	Used for upper and lower respiratory tract infections, whooping cough, diphtheria, or for other infections in patients who cannot take penicillins



By [kjaniskevich](https://www.cheatography.com/kjaniskevich/)

[cheatography.com/kjaniskevich/](https://www.cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 4 of 8.

Sponsored by [ApolloPad.com](https://www.apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://www.apollopad.com>

### Antibiotics - Classes of medication (cont)

<b>Aminoglycosides</b> <i>amikacin, gentamicin, streptomycin, tobramycin</i>	Inhibit bacterial protein synthesis and cause abnormal protein synthesis, Dose-dependent bactericidal Very effective, usually reserved for serious infections (like tuberculosis) or when other antibiotics have failed Require therapeutic drug monitoring – levels must be in specific range to be effective, but not toxic Injection or topical
<i>Gentamicin</i>	Used for serious (life-threatening) infections or when other antibiotics have failed (also topically as eye drops and creams/ointments – this would not require therapeutic drug monitoring)

### Antibiotics - Classes of medication (cont)

<b>Fluoroquinolones</b> <i>besifloxacin, ciprofloxacin, gatifloxacin, levofloxacin, moxifloxacin, norfloxacin, ofloxacin, floxacin</i>	Affect bacterial DNA synthesis, Bactericidal Most often used orally (also ear, eye) Absorption is affected by minerals (calcium, iron, magnesium) and need to be separated (~2h) Generally not used in children – affects cartilage development
<i>Ciprofloxacin</i>	Used commonly for respiratory, urinary, ophthalmic, gastrointestinal, and gynecological infections – high usage in community/out-patient
<b>Sulfonamides</b> <i>sulfacetamide, sulfamethoxazole, sulfapyridine</i> <i>Sulfa-</i>	Suppress bacterial growth by inhibiting essential folic acid needed within the cell, Bacteriostatic Broad spectrum, older class more resistance seen Orally and topically (acne) “Sulfa” is also a common “allergy”

### Antibiotics - Classes of medication (cont)

<i>Sulfamethoxazole-Trimethoprim (SMZ-TMP, Septra®, Bactrim®, -DS)</i>	Both drugs inhibit essential folic acid synthesis; work synergistically (a pharmacodynamic interaction) Used to treat urinary tract infections
<b>Carbapenems</b> <i>ertapenem, imipenem, meropenem, -penem</i>	Relatively new-ish Contain beta-lactam ring and inhibit cell wall synthesis (like penicillins) The beta-lactam ring is very resistant to destruction by penicillinase Broad spectrum – and very effective; as a newer class, they are being reserved for resistant infections (like MRSA, etc.)
<b>Miscellaneous VIPs</b>	
<i>Clindamycin</i>	protein synthesis inhibitor; bacteriostatic Used topically (acne), oral or IV for serious systemic infections High risk of superinfection (GI)

### Antibiotics - Classes of medication (cont)

<i>Nitro-fur-antoin</i>	inhibits protein, RNA, DNA, and cell wall synthesis; bactericidal Excreted unchanged through the kidney (no metabolism), therefore used only for urinary tract infections Changes urine to orange colour Must take with food
<i>Metro-nid-azole</i>	destroys bacterial DNA; bactericidal For anaerobic bacteria NO ALCOHOL USE (even small amounts present in cough syrup or mouthwash)

### Antibiotics - Classes of medication (cont)

<i>Vanco-mycin</i>	inhibits cell wall synthesis; bactericidal (through different mechanism than aminoglycosides) Reserved for severe infections that are resistant to anything else usually only used in hospital (methicillin-resistant staph. aureus - MRSA) Injection or oral If IV given too quickly Red Man Syndrome (flushing, red face, hypotension) slow down infusion Therapeutic drug monitoring required (like aminoglycosides)
<i>Linezolid</i>	inhibits bacterial protein synthesis Use to treat vancomycin-resistant enterococcus (VRE), pneumonia or skin infections caused by MRSA i.e. Severe infections resistant to other antibiotics

### Allergies- Classes of Medication

#### Antihistamines

### Allergies- Classes of Medication (cont)

1st-Generation Antihistamines	Block H1 receptors Shorter acting, cause more drowsiness, and work faster than 2nd Generation Used mostly to treat allergic response <i>Diphenhydramine</i> and <i>chlorpheniramine</i> most common Have anticholinergic effects Significant sedation – some use as a sleep aid
-------------------------------	--

#### 2nd-Generation

Antihistamines <i>Cetirizine (Reactine®)</i> , <i>loratidine (Claritin®)</i> , <i>desloratidine (Aerius®)</i> , <i>fexofenadine (Allegra®)</i>	<i>Cetirizine (Reactine®)</i> , <i>loratidine (Claritin®)</i> , <i>desloratidine (Aerius®)</i> , <i>fexofenadine (Allegra®)</i>
---	---

#### Intranasal Corticosteroids

To reduce inflammation in nasal mucous membranes, and local immunosuppression  
Used daily to prevent symptoms  
Can take up to 2 weeks for full effect  
Local administration prevents systemic side effects



By [kjaniskevich](https://www.cheatography.com/kjaniskevich/)

[cheatography.com/kjaniskevich/](https://www.cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 6 of 8.

Sponsored by [ApolloPad.com](https://www.apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://www.apollopad.com>

### Allergies- Classes of Medication (cont)

**Decongestants** Sympathomimetics – stimulants – cause vasoconstriction and reduction of mucous production  
*Phenylephrine, pseudoephedrine*  
 For immediate relief of nasal congestion – oral or intranasal  
 Short term-use only – rebound congestion if longer than 3-5 days (intranasal)

### Drugs for Anaphylaxis

**Epinephrine** Stimulates both  $\alpha$  and  $\beta$  adrenergic receptors  
 Via  $\alpha$ -receptors: counters the high vascular permeability that occurs during anaphylaxis that leads to loss of intravascular fluid and hypotension  
 Via  $\beta$ -receptors: causes bronchial smooth muscle relaxation and relieves bronchospasm, dyspnea, and wheezing  
 Also alleviates pruritus, urticaria, and angioedema

### Anti-fungals - Classes of medication

**Amphotericin B** Binds to fungal cell membranes, making them leaky, Given IV

**Azole Antifungals** Alter fungal cell membranes by depleting ergosterol  
*fluconazole, itraconazole, ketoconazole, miconazole, voriconazole*  
 Used orally, topically, injection; fluconazole available OTC  
 Safer than amphotericin B  
 Most often for vaginal candidiasis, athlete's foot, or thrush  
 metronidazole is NOT an azole antifungal

### Miscellaneous

*Ciclopirox* topical med used for fungal nail or scalp infections (nail polish or shampoo)

*Terbinafine* oral med for fungal nail infections

*Nystatin* cream available without prescription for many topical fungal infections (ringworm, diaper rash)

### Anti-Virals -Classes of medication

**HIV** Because of antiretroviral drugs, HIV patients are able to live symptom-free for much longer with very low counts of the retrovirus  
 Antiretroviral drugs block the HIV replication cycle

### Anti-Virals -Classes of medication (cont)

**HAART** – Goal is to reduce plasma HIV to its lowest possible level - highly active antiretroviral therapy  
 HIV still remains in the lymph nodes  
 Blood and lymph are separate rivers that cross occasionally  
 Use different classes of antiretrovirals at same time to reduce resistance  
 Each class 'attacks' different step of replication cycle

### Herpes Infections

HSV1: oral cold sores

HSV2: genital ulcerations

Zoster: shingles (due to previous varicella-zoster infection)

*Acyclovir, famciclovir, valacyclovir*  
 Mostly controlled by oral therapy of antivirals – taken at first sign of outbreak, continued for short term  
 These antivirals prevent viral DNA synthesis  
 Very well tolerated – take with food

### Over the Counter medications for Herpes

*Lipactin*® - can reduce pain, may speed healing  
 heparin + zinc  
 Mechanism does not match pathophysiology



By [kjaniskevich](https://kjaniskevich)

[cheatography.com/kjaniskevich/](https://cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 7 of 8.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>

### Anti-Virals -Classes of medication (cont)

*Abreva®* - prevents viral entry into cells, stops spread if caught early (can reduce duration of cold sore by ~1 day)

**Influenza** BEST PROTECTION = VACCINATION  
*Amantadine, and neuraminidase inhibitors (oseltamivir and zanamivir)* Antiviral drugs may decrease severity of symptoms of influenza and may shorten symptom time by a couple days IF taken within first 48 hours  
 Generally used only in patients at high risk of complications from influenza

### Immunosuppression - Classes of Medication

**Calcineurin Inhibitors** "Classic" immuno-suppressants used for transplants (or topically for psoriasis)  
*Cyclosporine, tacrolimus, pimecrolimus (topical)* Disrupt T-cell function by binding to calcineurin  
 They are not specific – suppress the ENTIRE immune system patient is very susceptible to any other infection  
 Extensive monitoring for detailed WBC counts and signs of infection (see slide on monitoring)

### Immunosuppression - Classes of Medication (cont)

**Corticosteroids** Anti-inflammatory and immunosuppressant activity  
 Often used to control exacerbations of condition such as asthma, rheumatoid arthritis, MS, etc.  
 Pulse therapy (very high doses, gradual taper) to minimize side effects  
 Many, many side effects

**Biologics** medications produced using biological processes in living organisms such as yeast and bacteria  
 Have active pharmaceutical ingredients that cannot reasonably be synthesized by chemical means (too complicated)  
 Are complex, large molecules derived from living sources and produced through a number of intricate steps  
 Biologics can be immunosuppressant or immunostimulant (very specifically) or replace a substance that is missing (insulin)

### Immunosuppression - Classes of Medication (cont)

Vaccines, Blood products, Hormones & growth factors, Enzymes Gene therapy, Cancer treatments

### Chemotherapy

*Cytotoxic drugs* traditional; interfere with or damage DNA, causing apoptosis (programmed cell death)

*Hormonal therapy* not cytotoxic; effects mediated through hormonal receptors (deprivation) – for hormone-responsive cancer (breast, prostate, etc.)

*Immunotherapy* monoclonal antibodies, vaccines; non-specifically boost immune system to help eradicate cancer (interferon alfa)

*Targeted agents* monoclonal antibodies, tyrosine-kinase inhibitors (TKIs); the future of treatment – to target cancer cells only

### Tuberculosis medications

*Rifampin (RMP)* Most potent anti-TB drug available  
 Good bactericidal activity, prevents acquired drug resistance and is very important in preventing relapse  
 Current doses are based on studies performed in the 1960s, when the lowest effective dose was used because of the high cost of the drug; concerns now that dose is too low -> current trials -> dosing recommendations may change



By [kjaniskevich](https://kjaniskevich)

[cheatography.com/kjaniskevich/](https://cheatography.com/kjaniskevich/)

Published 13th April, 2021.

Last updated 13th April, 2021.

Page 8 of 8.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>