

Infective Endocarditis

Causative organisms

Staph. aureus, group D strep, enterococci, HACEKs

Organism in IVDA

Staph. aureus (tricuspid valve*)

Organisms prosthetic valve endocarditis

Staph, gram-, fungi (first 2 months) and staph/strep after that

Regurgant valve defect

Seen in most endocarditis pts, makes them more susceptible

How infections occur

Direct intravascular contamination or from bacteremia from surgeries

Classic features

Osler nodes, Janewar lesions, Roth spots, petechiae, splinter hemorrhages

Duke Criteria

Used to establish diagnosis

Treatment

Vancomycin + Ceftriaxone

Indications for Abx Prophylaxis

If pts. with prosthetic valves, congenital heart disease, valve disorder, transplants are going to get dental work or surgery

Prophylactic abx

Amoxicillin

Aortic Aneurysms

Definition

Weakness and subsequent dilation of the vessel wall, usually from a genetic defect or atherosclerotic damage to the intima

Most common

Atherosclerosis (can see in Marfan's or Ehlers-Danlos though)

Classic clinical scenario

Elderly male smoker with CAD, emphysema, and renal impairment

Where are they found

90% abdominal, 10% thoracic

Clinical features

Pulsatile abdominal mass +/- abdominal or back pain

Symptoms of AAA

Severe back, abdominal, or flank pain. Hypotension + rupture shock

Lab Studies

Abdominal U/S followed by CT w/ contrast

Treatment

Endovascular or open surgical repair

Giant Cell Arteritis

Definition

Systemic inflammatory condition of medium & large vessels, pts. >50yo, often coexists with PMR

Most commonly-affected arteries

Temporal artery

Consequence of not treating aggressively

Blindness

Clinical Features

Headache, scalp tenderness, jaw claudication, throat pain, visual abnormalities

Lab Studies

ESR + CRP both elevated

Definitive diagnostic

Temporal artery bx

Treatment

High-dose prednisone x few months + ASA

Rheumatic Heart Disease

Rheumatic Fever

A systemic immune response occurring 2-3 weeks after a Beta-hemolytic strep. pharyngitis

Valve most commonly involved

Mitral valve (75-80%), then aortic valve (30%)

Jones Criteria

Diagnostic criteria to establish diagnosis

Treatment

Bedrest, salicylates, IM penicillin, and early treatment of strep pharyngitis for prevention*



By **ksellybelly**
cheatography.com/ksellybelly/

Published 24th July, 2014.
Last updated 13th May, 2016.
Page 1 of 2.

Sponsored by **CrosswordCheats.com**
Learn to solve cryptic crosswords!
<http://crosswordcheats.com>

PVD—Chronic Venous Insufficiency

Definition

Loss of wall tension in veins, resulting in stasis of venous blood and often associated with a history of DVT, leg injury, or varicose veins

Clinical Features

Progressive edema starting at ankle, skin changes, itching, dull pain with standing and ulceration, skin is shiny/thin/atrophic with dark pigmentary change and subcutaneous induration, stasis ulcers above ankle

General Treatment

General: leg elevation, avoidance of sitting/standing, compression hose.

Treatment for Stasis Dermatitis

Wet compresses, HC cream, Zinc oxide, anti fungal cream (ulcerations may need graft)

PVD—Peripheral arterial disease

Etiology

Atherosclerosis or thromboembolism (trauma, hyper coagulable states, etc.)

Clinical features

lower leg pain with exercise which is relieved by rest (AKA intermittent claudication), progresses later to pain at rest, numbness, tingling, ischemic ulcerations, gangrene

The "Ps" of extremity occlusion

Pain, pallor, pulselessness, paresthesias, paralysis, poikilothermia

Lab studies/diagnostics

Doppler flow studies, ABI,

Treatment

Cilostazol + antiplatelet rx + lifestyle (NO smoking, more exercise), surgery and revascularization

PVD—Thrombophlebitis & DVT (cont)

Features of superficial thrombophlebitis

Dull pain, erythema, tenderness, induration. Most common in long saphenous vein.

Class findings of DVT

Swelling of the involved area and redness

Diagnostic Studies

Duplex U/S

D-Dimer

Highly sensitive, if <500 then negative, can rule out DVT

Treatment

Anticoagulation with LMWH (Lovenox), or heparin then warfarin

PVD—Varicose veins

Etiology

Superficial venous insufficiency and valvular incompetence

Clinical features

Dilated, tortuous veins, esp. long saphenous vein

Treatment

Compression stockings, leg elevation, exercise, laser ablation, endovenous radiofrequency, compression sclerotherapy

PVD—Thrombophlebitis & DVT

Thrombophlebitis

Involves occlusion of a vein + inflammatory changes

Virchow's Triad

Stasis + vascular injury + hyper-coagulability (predispose veins)

Most common place to find a DVT

Lower extremities and pelvis

Risk factors for DVT

Major surgery (total hip), long plane ride, hormone/contraceptive therapy, prolonged bed rest



By **ksellybelly**
cheatography.com/ksellybelly/

Published 24th July, 2014.
Last updated 13th May, 2016.
Page 2 of 2.

Sponsored by **CrosswordCheats.com**
Learn to solve cryptic crosswords!
<http://crosswordcheats.com>