

### Test 2 Q&A

True or False :

Q1) The probability which comes from the fact that you believe that you have a 1/2 chance of drawing a red card from a standard deck of cards is empirical probability. FALSE

Q) The events "not shutting off the water in the kitchen sink" and "receiving a high water bill" are not independent. TRUE

Q) The events "choosing a student who is male or a finance major from a nearby university to participate in a research study", are mutually exclusive. FALSE

Q)  $8! = 20160$ . False

Q2) Suppose the local shop makes dining room tables in three shapes and in three different finishes. Their production of the last six months is summarized in the following table:

Suppose one table is selected at random, Find the following:

A)  $P(\text{the table is oval}) = 177/400$  B)  $P(\text{the table is pine or rectangular}) = 101+47+26/400 = 177/400$  C)  $P(\text{the table is round and mahogany}) = 43/400$  D)  $P(\text{the table is oak} \mid \text{it is round}) = 52/129$  E)  $P(\text{the table is oval} \mid \text{it is mahogany}) = 62/152$ .

Q3) A coordinator will select 7 songs from a list of 13 songs to compose an event's musical entertainment lineup. How many different lineups are possible? A:  ${}^{13}P_7 = 8648640$

Q6) A box contains 16 white marbles and 12 green marbles. If two are drawn without replacement, what is the probability that they are both green?  $12/28 * 11/27 = 132/756$ .

Q7) The local restaurant is offering a dinner special that includes an appetizer, an entree, and a dessert. If there are 5 different appetizers to choose from, 11 different entrees, and 4 different desserts, how many different specials are possible?  $5 * 11 * 4 = 220$

### info

### info (cont)

> The median is the 50th percentile; the point in data where 50% of the data falls below that point and 50% fall above it. Steps to calculate the K percentile  
**1)** Order all the values in the data set from smallest to largest.  
**2)** Multiply K% by the total number of values, n(index).  
**3)** IF the index or n is not a whole number, round up to the nearest whole number. Count the values in your data set from left to right (smallest -> largest) until you reach the index. The corresponding value is your K %ile.  
**4)** Count the values in your data set from the left until you reach the index. The K %ile is the average of that corresponding value in your data set & the value that directly follows it.

IE: 25 test scores in order

{43,54,56,61,62,66,68,69,70,71,72,77,78,79,85,87,88,89,93,95,96,98,99,99} To find the 90th %ile  
**1)** Multiply  $0.90 * 25(\text{amount of data}) = 22.5(\text{index})$ .  
**2)** Rounding up to you get 23, so counting from L->R until you reach the 23rd value (98) that is the 90th %ile of this data set. OR if you wanted the 20th %ile  
**1)**  $0.20 * 25 = 5(\text{index})$ .  
**2)**  $5\text{th} + 6\text{th value} = (62+66)/2 = 64$  is the 20th %ile. The median is the 13th score (77).

### BASIC PROBABILITY

**Probability Experiment** - Any process with a result determined by chance.

**Sample Space** - Each individual result that's possible for a probability experiment.

**Event** - A subset of outcomes from the sample space.

IE: Consider the experiment of rolling a dice. There are 6 possible outcomes, namely the number 1-6. The **sample space** is the set of all outcomes which in this case is simply {1,2,3,4,5,6}. The **event** "rolling an even number" is the subset of outcomes {2,4,6}. On the other hand, the event "rolling a number less than 10" is the set {1,2,3,4,5,6}.

**Subjective Probability** - The least precise type of probability.

**Experimental Probability** - specifically it is calculated by dividing the number of times an event occurs by the total number of trials performed. F: If E is an event, the **P(E)**, "the probability that E occurs is given by:  $P(E) = f/n$  F= the frequency of event E, N= the total number of times the experiment is performed. IE: A fisherman wants to know the prob that a fish he catches in his fav pond will be a catfish. HE records the numbers & types of fish he catches for one week. He catches 92 fish, 43 of which were catfish. He then calculates the prob as follows:  $\rightarrow P(\text{catfish}) = \# \text{ of catfish caught} / \text{total} \# \text{ of fish caught} \rightarrow 43/92 = 0.4674$ .  
**Law of Large Numbers** - The greater the number of trials, the closer the experimental probability will be to the true prob.

### CLASSICAL PROBABILITY

The most precise type of probability. It is calculated by taking all possible outcomes into consideration. It states that if all outcomes are equally likely; the probability of an event is equal to the number of outcomes included in the event divided by the total number of outcomes in the sample space.

$P(E) = n(E)/n(S)$   $n(E)$  = the number of outcomes in the event  $n(S)$  = number of outcomes in the sample space.

**Empirical Rule (68-95-99.7%)** - Use the empirical rule to estimate the probability of a porcupine living between 20.5 & 27.4 years with an average lifespan of 22.8, the standard deviation is 2.3 years. Chart 1.0 **Step 1)** The average lifespan 22.8(mean). **2)** SD is 2.3 years so 1 SD below is 20.5 & 1 above is 25.1 yrs. **3)** Two SD's below the mean is 18.2, two above is 27.4. **4)** Three SDs below is 15.9yrs & above is 29.7yrs. **5)** Looking for prob of porcupine living between 20.5-27.4 years. **6)** Empirical Rule tells us 95% of the porcupines will have lifespans within 2 SD's of the avg. Also 68% will have lifespans within 1 SD of the mean. **7)** That leaves 95% - 68% = 27% of porcupines between 1&2 SD's or 13.5%(each side). **8)** The probability is 68% + 13.5% or 81.5%

#### **STANDARD SCORE (Z-SCORE)**

A z-score is defined as the number of standard deviations a specific point is away from the mean.

Q) The grades on a physics midterm at Santa Rita are normally distributed with  $M = 78$  &  $\sigma = 3.5$ . Ishaan scored 79% on the exam. Find the Z-Score  
Steps 1)  $Z = \frac{79 - 78}{2.5} \rightarrow Z = 0.29$ . Ishaan's score was 0.29 SD's above the mean.

#### **PERCENTILE**

**IE:1** All fish from the pond recorded, total # of fish 1235, 541 are catfish. So:

$$P(\text{catfish}) = \frac{n(E)}{n(S)} \rightarrow \frac{541}{1235} = 0.4381$$

**IE:2** Idea of rolling a 6 sided die and getting an even number. There are 3 possible outcomes a 2,4, or 6. So  $n(E) = 3$  with 6 outcomes in the sample space thus  $n(S) = 6$ . **P(even) =  $\frac{3}{6} \rightarrow \frac{1}{2} = 0.5$  prob**

**Subjective Probability-** An educated guess regarding the chance that an event will occur.



By **MADdogz43**

[cheatography.com/maddogz43/](https://cheatography.com/maddogz43/)

Published 18th March, 2015.  
Last updated 13th May, 2016.  
Page 1 of 4.

Sponsored by **ApolloPad.com**  
Everyone has a novel in them. Finish Yours!  
<https://apollopad.com>



= 4/51 So -->

$P(\text{king \& queen, w/o replacement}) = P(\text{king}) * P(\text{queen, given king drawn first})$

= 4/52 \* 4/51 = 1/13 \* 4/51 = 4/663 = 0.0060

**Conditional Probability** Denoted  $P(F | E)$ , it is the probability of event F occurring given that event E occurs first.  $P(F | E) = P(E \& F) / P(E)$

**IE:** One card has already been chosen, w/o replacement. What is the probability of now choosing a second card from the deck and it being red, given that the first card was a diamond? Given that the first card was a diamond (red card) there is only 25 red cards left instead of 26, and only 51 total cards remaining. So :  $P(\text{red} | \text{diamond}) = 25/51 = 0.4902$

**Multiplication Rule for Probability of Dependent Events**

$P(E \& F) = P(E) * P(F | E)$



By MADdogz43

Published 18th March, 2015.  
Last updated 13th May, 2016.  
Page 2 of 4.

[cheatography.com/maddogz43/](http://cheatography.com/maddogz43/)

Sponsored by [ApolloPad.com](http://ApolloPad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>