

### Number of protons, neutrons and electrons

Number of protons = atomic number

Number of neutrons = mass number - atomic number

Number of electrons = atomic number - the charge

Example:

Carbon - 12 has an atomic number of 6 and a mass number of 12

number of protons = 6

number of neutrons =  $12 - 6 = 6$

number of electrons =  $6 - 0$  (neutral atom) = 6

if it is an isotope/has a charge:

Carbon - 13

number of protons = 6

number of neutrons =  $13 - 6 = 7$

number of electrons =  $6 - +1 = 5$  (C-13 has a +1 charge, so it loses 1 electron)

### Naming transition metals

For binary compounds (those with only two elements), the naming convention follows a specific set of rules.

However, compounds involving a metal and a non-metal, like iron chloride, the non-metal is typically ended with an 'ide'.

Non-metals typically form a negatively charged ion (anion), which are named with an 'ide'.

This is why both  $\text{FeCl}_2$  and  $\text{FeCl}_3$  are referred to as "chloride."

Chlorine forms the chloride ion when it gains an electron, becoming  $\text{Cl}^-$ .

In the case of iron chloride, the 'ide' ending used in both  $\text{FeCl}_2$  and  $\text{FeCl}_3$  doesn't refer to the number of chlorine present, but the nature of the charge formed by the chlorine, which is an anion

The use of 'ate' and 'ite' are typically used when the non-metal in a compound is oxygen

- Chlorate ( $\text{ClO}_3^-$ ) and chlorite ( $\text{ClO}_2^-$ ) have suffixes indicating the number of oxygen present/the specific arrangement of atoms around oxygen

This naming convention doesn't directly apply to compounds involving chlorine and other elements like iron.

Instead, they are named according to their oxidation states. When an element can have multiple oxidation states, it's indicated using Roman numerals.

So, to name transition metals, you must first figure out the oxidation state of the compound, and indicate this using roman numerals

Example:



- The iron here has a 2+ oxidation state, meaning it's lost 2 electrons

- Therefore, it's named Iron (ii) Chloride



- In this case, iron has lost 3 electrons making it +3



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 1 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Naming transition metals (cont)

- So it's named Iron (iii) chloride

Another example CuS - Copper sulfide

To figure out if it's i-iii, we figure out the charge of copper by first figuring out the charge of sulfur

- Sulfur is a chalcogen found in group 6a, and they form  $2^-$  anions

- Copper can form either +1 or +2 ions, so in this case it would have to be a +2 charge to balance out the charge of sulfur.

*- if the charge of copper was +1, then 2 copper atoms would be needed to balance the compound, but given that there is only 1 in CuS, we can determine that the charge would be +2*

Since CuS is a neutral compound, the total positive charge of copper must balance the total negative charge of sulfur by determining copper's charge, which we already determined was -2. since the compound is neutral, the total sum of charges must = 0

We denote the charge of copper as 'x'

-  $x + (-2) = 0$

We denote the charge of copper as "x." Since the compound is neutral, the sum of the charges of copper and sulfur must equal zero:

-  $x + (-2) = 0$

- solving for x, we find that  $x = +2$ , meaning copper must have a +2 charge

Therefore, in naming CuS, we put the number of the charge in roman numerals

- CuS = Copper(ii) sulfide.

In the case of Cu<sub>2</sub>S, we'd follow the same steps

- S has a charge of -2, we must balance the copper

-  $2x + (-2) = 0$

- solving for x, we find  $x = +1$  because  $1 \times +1 + (-2) = 0$

Therefore, copper has a +1 charge

so Cu<sub>2</sub>S = Copper (i) sulfide

### Average atomic mass of isotopes

Atomic mass x abundance of isotope for each isotope, then add together

Example:

Iron has 4 isotopes.

Fe-54, 56, 57, and 58

To calculate the average atomic mass, first get the abundance % and atomic mass of each isotope.

Fe-54

abundance % = 5.845%

Atomic mass = 53.9396

Convert abundance to decimal format by dividing by 100  
= 0.05845

Now, multiply the atomic mass with the % as a decimal

$5.845 \times 0.05845 = 0.34164025$

Do the same for the remaining isotopes, then add the final numbers together

= 55.845

Highest abundance % is the most commonly found isotope btw

### max obtainable mass

Maximum mass that can be obtained

Laughing gas" or nitrous oxide, N<sub>2</sub>O, is prepared by the thermal decomposition of ammonium nitrate:  $\text{NH}_4\text{NO}_3 (\text{s}) \rightarrow \text{N}_2\text{O} (\text{g}) + 2 \text{H}_2\text{O} (\text{l})$  (a) What is the maximum mass in grams of N<sub>2</sub>O (g) that can be obtained from 1.53 kg of ammonium nitrate?

Calculate moles of ammonium nitrate

- Mass is 153g, molar mass is 80g/mol, divide mass by molar mass

- Moles =  $153/80 = 1.91$  moles

According to equation, 1 mole of NH<sub>4</sub>NO<sub>3</sub> yields 1 mole of N<sub>2</sub>O

Number of moles of N<sub>2</sub>O produced will also be 1.91

Convert moles of N<sub>2</sub>O to grams

Multiply N<sub>2</sub>O molar mass (44.02) by number of moles

Mass of N<sub>2</sub>O =  $1.91 \times 44.02\text{g/mol} = 84.1$  grams

If the percentage yield was 76%, what mass (grams) of N<sub>2</sub>O was actually produced?

Calculate maximum theoretical yield – 84.1 grams

Apply percentage yield to find actual yield

### max obtainable mass (cont)

Given the percentage yield as 76%, we multiply it by the maximum theoretical yield

Actual yield =  $84.1 \text{ g} \times 76/100 = 63.9\text{g}$

### % (w/w)

Percentage by weight, indicating the mass of the solute per 100 grams of total product (solution)

Tells us the proportion of the solute to the entire product

Number of grams of NaOH present in a 375g can of oven cleaner labelled 4.2% (w/w) NaOH

Calculate mass of NaOH per gram

Given percentage of NaOH in oven cleaner is 4.2% w/w

To convert % to grams/gram, divide it by 100

-  $4.2/100 = 0.042\text{g}$

Determine total mass of NaOH in the can

Total mass is given as 375g

To find mass of NaOH in the can, you multiply the mass of NaOH per gram of product by the total mass of the can

- NaOH mass in can =  $0.042 \text{ g/g} \times 375\text{g}$

- = 15.75g

### Evaporation/ %Mass

Calculate concentration of a solid as a percentage by mass (% mass)

% of mass calculated by dividing the mass of the solute by the total mass of the solution and then multiply by 100

Calculate the concentration of the solid mass per unit volume g/L

This concentration represents the amount of solid material (solute) per unit volume of the solution

A 1.62 kg (1.71 L) sample of creek water was evaporated to dryness, leaving 6.33 g of solid material.

Calculate the concentration of this solid as percentage by mass and calculate the concentration of this solid mass per unit volume g/l

Concentration of the solid as a percentage by mass (% mass)

The mass of the solid material left after evaporating is 6.33g

The total mass of the solution is 1.62kg = 1620g



By massiveballs

Not published yet.

Last updated 12th April, 2024.

Page 3 of 18.

Sponsored by ApolloPad.com

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Evaporation/ %Mass (cont)

% mass =  $(6.33\text{g}/1620\text{g}) \times 100\%$  - 0.391%  
 Calculate the concentration  
 The volume of creek water is 1.71L  
 Concentration = mass of solute / volume of solution  
 =  $6.33\text{g}/1.71\text{L} = 3.70 \text{ g/L}$

### concentration in dissolved solution

Figure out number of moles first  
 Molarity = moles per litre ( $\text{mol}\cdot\text{L}^{-1}$ )  
 work out number of moles first, then divide by the volume  
 Molar mass of Na = 22.99  
 MM of Cl = 35.45  
 divide by the given volume (6g)  
 $6/(22.99 + 35.35 \text{ moles})$   
 =  $6/58.44 \text{ mol}$   
 =  $0.103 \text{ gmol}^{-1}$   
 volume in mL, convert to L  
 $750\text{ml}/1000 = 0.75\text{L}$   
 concentration is the mols/volume  
 $0.103/0.75$   
 = 0.14M


### Moles corresponding to molecules

convert number of molecules to moles by dividing given number of molecules by Avo number  
 Number of moles = number of molecules/ $6.022 \times 10^{23}$   
 Moles of CH<sub>4</sub> corresponding to  $1.56 \times 10^{20}$  molecules  
 $N(\text{CH}_4) = 1.56 \times 10^{20} \text{ molecules}/6.022 \times 10^{23}$   
 =  $25.9 \times 10^{-4}$

### Mass of product

**If 100.0 g of Al is added to 0.11 mol of O<sub>2</sub>, how many grams of Al<sub>2</sub>O<sub>3</sub> (s) will be produced?**

Write balanced equation



- 4 moles of Al react with 3 moles of oxygen to make 2 moles of aluminum oxide

Moles of Al using its molar mass

= moles of Al = mass of Al/Molar mass of Al

=  $100\text{g}/27 \text{ g/mol} = 3.70 \text{ mol}$

Mols of oxygen are already given, they are 0.11 mol

Determine limiting reactants

we assume 1 is in excess

- We use stoichiometry to determine how much of the O<sub>2</sub> would be needed to react completely with the excess reactant. if there is more of the other reactant than the calculated amount, then it is in excess; otherwise it is the limiting reactant.

- to react all aluminium, it would require  $3/4 \times$  moles of Al

-  $3/4 \times 3.70 = 2.78 \text{ mol of O}_2$

Assuming O<sub>2</sub> is in excess, it would require  $4/3 \times$  moles of O<sub>2</sub>

=  $4/3 \times 0.11 = 0.15 \text{ mol Al}$

Since there's less than 2.78 mol O<sub>2</sub> to react with the Al, and more than 0.15 mol Al to react with the O<sub>2</sub>, O<sub>2</sub> is the limiting reactant

Calculate moles of Al<sub>2</sub>O<sub>3</sub>

from the equation, we see that 4 moles of Al react with 3 moles of O to produce 2 moles of Al<sub>2</sub>O<sub>3</sub>.

Since O is limiting reactant, we use its moles to find the moles of Al<sub>2</sub>O<sub>3</sub>

moles of Al<sub>2</sub>O<sub>3</sub> =  $2/3$  moles of O<sub>2</sub>

=  $2/3 \times 0.11 = 0.073$

Mass of Al<sub>2</sub>O<sub>3</sub> produces

= Moles of Al<sub>2</sub>O<sub>3</sub> x molar mass of Al<sub>2</sub>O<sub>3</sub>

=  $0.073 \times 102.0 = 7.48 \text{ grams}$



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 4 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### lewis structure

Count total number of valence electrons and add them together

- for example,  $\text{NCl}_3$ , nitrogen has 5, chlorine has 7. since there are 3 Cl atoms, that's  $3 \times 7 = 21$  valence electrons.
- $5 + 21 = 26$  valence electrons for the entire molecules

Choose the central atom

- the central atom is the less electronegative one, as it can make more bonds
- $\text{NCl}_3$ , nitrogen is the central atom

Connect the atoms with single bonds

- Connect the central nitrogen atom to each of the 3 chlorine atoms using single bonds
- this uses up 3 pairs of electrons,  $26 - 3 = 23$  electrons remaining

distribute the remaining electrons

- place lone pairs on the outer atoms first, and then fill the remaining electrons around the central atoms
- each lone pair is represented by 2 electrons

check for formal charges after

- a formal charge occurs when an atom doesn't have the expected number of valence electrons
- calculate the formal charge for each atom by subtracting the number of lone pair electrons and half the number of bonding electrons from the number of valence electrons the atom brings

If any atoms have formal charges, try to minimize them by moving lone pairs or changing the arrangement of bonds

- the goal is to have the lowest formal charges possible while still satisfying the octet rule

N - 1 lone pair (2 electrons)  
 Each Cl - 3 lone pairs (6 electrons)

### mass of production in reaction

$\text{KCN (aq)} + \text{HCl (aq)} \rightarrow \text{KCl (aq)} + \text{HCN (g)}$

**If 0.250 g KCN reacts with excess acid, calculate the mass (in g) of HCN produced?**

Determine molar masses of the substances involved

- Molar mass of KCN = molar mass of K + molar mass of C + molar mass of N
- Molar mass of HCN = molar mass of H + C + N

convert the mass of KCN to moles

- number of moles = mass/molar mass

### mass of production in reaction (cont)

apply stoichiometry to find the molar ratio of KCN and HCN to find the number of moles of HCN produced when a certain number of moles of KCN reacts

Convert moles of HCN to mass

- mass = number of moles x molar mass

Moles of KCN =  $0.250/65$   
 $= 0.003846$

the ratio is 1:1, so the number of HCN moles is the same as the number of KCN moles

To find the mass of HCN, multiply the number of moles of HCN by its molar mass

Moles of HCN x Molar mass of HCN  
 $= 0.003846 \text{ mol} \times 27\text{g/mol}$   
 $= 0.104 \text{ g}$

mass of HCN produced when 0.250 g of KCN reacts with excess acid is 0.104

### concentration of a solution

**"Calculate the concentration (mol/L) of a saline solution of 9 g table salt in 1 L water."**

Determine the molar mass of NaCl

- sum of the atomic mass of Na and Cl
- $\text{MM Na} + \text{MM Cl}$
- $22.99 \text{ g/mol} + 35.45 \text{ g/mol}$
- $58.44 \text{ g/mol}$

Convert the mass to moles

- the given mass of table salt is 9g
- number of moles of NaCl
- $\text{Mass of NaCl} / \text{molar mass of NaCl}$
- $9\text{g} / 58.44\text{g/mol}$
- $0.154 \text{ mol}$

Now calculate the molarity/concentration of the NaCl in the saline solution

Concentration (mol/L) = number of mols of solute/ volume of solution (L)

- $0.154\text{mol} / 1\text{L}$
- $0.154 \text{ mol/L}$

The concentration of the saline solution, 9g of NaCl in 1L of water, is 0.154 mol/L



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 5 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>

### concentration of a solution (cont)

**Explain how you would proceed to make 50 ml of 0.030 M solution out of the above stock saline solution.**

Calculate volume of stock solution needed

use the dilution formula to calculate the volume of the stock solution (0.154 M) to prepare the desired result:

$$C_1V_1 = C_2V_2$$

-  $C_1$  = concentration of stock solution (0.514 M)

-  $V_1$  = volume of stock needed (?)

-  $C_2$  = desired concentration of final solution (0.030 M)

-  $V_2$  = Volume of final solution (50mL)

Solve for  $V_1$

$$V_1 = C_2 \times V_2 / C_1$$

$$= (0.030 \text{ mol/L}) \times (0.050 \text{ L}) / 0.514 \text{ mol/L}$$

$$= 0.0015 / 0.154 \text{ L}$$

convert to mL

$$= 0.0097 \times 1000 \text{ mL/L}$$

$$= V_1 = 9.7 \text{ mL}$$

You need 9.7 mL of the solution to prepare 50 mL of the desired solution.

### water solubility and ions

Determining which compounds will form ions when dissolved in water.

Identify if compound is ionic, polar covalent or non-polar covalent

- ionic: consist of metal and a nonmetal or a metal and a polyatomic ion

- polar: significant different in electronegativity btwn bonded atoms, resulting in partial pos/neg charges

- non polar: similar electronegativity, resulting in equal sharing of electrons and no significant charge separations

Solubility rules:

- Most nitrate ( $\text{NO}_3^-$ ), acetate ( $\text{C}_2\text{H}_3\text{O}_2^-$ ), and chlorate ( $\text{ClO}_3^-$ ) salts are soluble

- most alkali metal (group 1) and ammonium salts ( $\text{NH}_4^+$ ) salts are soluble

- most chloride ( $\text{Cl}^-$ ), bromide ( $\text{Br}^-$ ), and iodide ( $\text{I}^-$ ) salts are soluble, except for those of silver, lead(ii) and mercury(i)

### water solubility and ions (cont)

- most sulfate ( $\text{SO}_4^{2-}$ ) salts are soluble, except for those of calcium, strontium, barium, lead(ii), and some silver salts

#### Dissociation in water

ionic compounds with ions are soluble according to the rules, it will dissociate into ions when dissolved in water

if polar or nonpolar, it wont

Some compounds that fully dissociate into ions in solution are considered strong electrolytes

- these compounds conduct electricity well in solution due to the presence of free ions

- strong acids, bases and soluble ionic compounds

Some dissociation behaviour of a compound can vary, some partially dissociate into ions while others fully dissociate

- weak acids and bases partially dissociate into ions in solution, theyre weak electrolytes

- non-electrolytes dont dissociate into ions when dissolved in water

#### Example




Lithium sulfate, containing lithium (Li) and sulfate ( $\text{SO}_4^{2-}$ ) ions.

Li salts, such as lithium sulfate, are generally soluble in water as lithium compounds are highly soluble

most sulfate salts are soluble

based on this, lithium sulfate will dissociate into its constituent ions

-  $\text{Li}^+$  and  $\text{SO}_4^{2-}$



### volume of acid to neutralise

volume 0.355 M perchloric acid to neutralise 15.5mL of a 0.179 calcium hydroxide solution

Write a balanced equation for the reaction

- perchloric acid ( $\text{HClO}_4$ ) and calcium hydroxide ( $\text{Ca(OH)}_2$ )



Determine the stoichiometry

From the equation, you can see that 2 moles of  $\text{HClO}_4$  react with 1 mole of  $\text{Ca(OH)}_2$ , meaning the stoichiometric ratio is 2:1

Calculate the moles of  $\text{Ca(OH)}_2$



By massiveballs

Not published yet.

Last updated 12th April, 2024.

Page 6 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### volume of acid to neutralise (cont)

Use the given concentration and volume of the calcium hydroxide solution to calculate the number of moles present.

- Moles of  $\text{Ca}(\text{OH})_2 = \text{concentration} \times \text{volume}$

$$= 0.179 \text{ M} \times 15.5 \text{ mL} / 1000 \text{ L}$$

$$= 0.179 \text{ M} \times 0.0155 \text{ L}$$

$$= 0.00277 \text{ Mol}$$

find the moles of  $\text{HClO}_4$

since the ratio between them is 2:1, the number of moles of perchloric acid needed is twice the number of calcium hydroxide

- moles of  $\text{HClO}_4$

$$= 2 \times 0.00277$$

$$= 0.00554 \text{ Mol}$$

now use the concentration of perchloric acid solution to find the volume of it thats needed

Volume = moles/concentration

$$= 0.00554 \text{ mol} / 0.355 \text{ M}$$

$$= 0.00554 / 0.355 \text{ L}$$

$$= 0.0156 \text{ litres}$$

So approx. 0.0156 litres or 15.6 mL of 0.355 M of perchloric acid needed to neutralise 15.5mL of 0.179 M calcium hydroxide solution

### Molecular formula from % composition

Follow previous steps to get the empirical formula, then find the molar mass of it by adding up the atomic masses of all atoms in the formula

Determine the molecular formula mass

You need to know the molecular weight/molar mass of the compound, if not known, you have to determine it experimentally

Calculate the 'multiplier' or 'factor'

Divide the molecular weight of the compound by the empirical formula mass to get the 'multiplier' that represents how many times the empirical formula must be multiplied to get the molecular formula

Multiply the subscripts in the empirical formula by the 'multiplier'

**Example:**

### Molecular formula from % composition (cont)

Compound: 40% carbon, 6.7% hydrogen, and 53.3% oxygen, with a molecular weight of 180 g/mol

the empirical formula is  $\text{CH}_2\text{O}$

to find the mass of it:

Mass of C = 12.01 g/mol

Mass of H = 1.01 g/mol

Mass of O = 16 g/mol

The empirical formula mass:

$$(12.01 \times 1) + (1.01 \times 2) + (16.00 \times 1)$$

$$= 30.03 \text{ g/mol}$$

Calculate the 'multiplier'.

$$\text{- Multiplier} = \text{Molecular weight} / \text{Empirical formula mass} = 180 / 30.03 = 6$$

Multiply the subscripts in the empirical formula by the multiplier

$$\text{C}_1\text{H}_2\text{O}_1 \times 6 = \text{C}_6\text{H}_{12}\text{O}_6$$

so the molecular formula is  $\text{C}_6\text{H}_{12}\text{O}_6$

By following these steps, you can determine the molecular formula of a compound from its percent composition and molecular weight.

### Excess reactant

Write balanced equation and determine limiting reactant

- calculate moles of each reactant involved and compare these values to determine which is limiting and which is in excess

- the limiting reactant is the one that produces the least amount of product

Calculate theoretical yield

use the stoichiometry of the balanced equation

- multiply the number of moles of the limiting reactant by the stoichiometric coefficient of the product in the balanced equation to get the theoretical yield in moles.

Determine the excess reactant

This is the one that is not limiting

- subtract the amount of excess reactant that reacts from the initial amount of excess reactant given

now calculate the amount left after the reaction



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 7 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>


### Excess reactant (cont)

- subtract the amount of excess reactant that reacted from the initial amount of excess reactant given.

Example:

Hydrogen gas and oxygen gas to form water

balance equation:



- Suppose we have 10.0 grams of hydrogen gas and 20.0 grams of oxygen gas.

Determine limiting reactant

calculate moles of each reactant

- H<sub>2</sub> molar mass = 2.02g/mol

- O<sub>2</sub> molar mass = 32 g/mol

divide by the respective grams amount

- H<sub>2</sub> - 10.0g / 2.02g/mol = 4.95

- O<sub>2</sub> - 20.9g / 32g/mol = 0.625

O<sub>2</sub> has fewer moles, so its the limiting reactant, while H<sub>2</sub> is the excess

Calculate theoretical yield

Balanced equation shows us that 2 moles of H<sub>2</sub> react with 1 mole of O<sub>2</sub> to produce 2 moles of H<sub>2</sub>O.

O<sub>2</sub> is limiting, so the amount of H<sub>2</sub>O produced is determined by its quantity

- the theoretical yield of H<sub>2</sub>O is 2 x 0.625 mol = 1.25 mol

Since all the O<sub>2</sub> is consumed, we don't need to calculate the amount of excess reactant (H<sub>2</sub>) that reacts.

Since all O<sub>2</sub> was consumed, and the initial moles of H<sub>2</sub> was 4.95, all moles of H<sub>2</sub> will react, leaving no excess H<sub>2</sub> after reaction.

**7g ammonia reacted with 10 g of oxygen, which reagent is in excess?**

Determine number of moles of each reactant

$N = \text{mass (gs)}/\text{molar mass}$

NH<sub>3</sub>/ammonia

$N(\text{NH}_3) = 7\text{g}/17.034\text{g/mol}$

For O<sub>2</sub>/oxygen

$N(\text{O}_2) = 10\text{g}/32\text{g/mol}$

Compare the number of moles to find the limiting reagent

### Excess reactant (cont)

Use the stoichiometric ratio 4:5 moles of NH<sub>3</sub> react with 1 mole of O<sub>2</sub>

$N(\text{NH}_3) = 4/5 \times n(\text{O}_2)$

Calculate the number of moles of NH<sub>3</sub> using the same ratio

Identify the limiting reagent by comparing the calculated moles of NH<sub>3</sub> and O<sub>2</sub>

If  $n(\text{NH}_3) < n(\text{O}_2)$ , NH<sub>3</sub> is limiting, and vice versa

O<sub>2</sub> is the limiting reagent, bc  $n(\text{O}_2) = 0.411$  moles, which is less than  $n(\text{NH}_3) = 0.25$  moles

Calculate the mass of the product based on the limiting reagent

Use the stoichiometry to find the number of moles of the product NO, then use the molar mass of NO to find the mass of the product

$\text{Mass}(\text{NO}) = n(\text{NO}) \times \text{molar mass}(\text{NO})$

$0.25 \text{ moles} \times 30\text{g/mol}$

$= 7.5\text{g}$ , round to 8g

### Moles of substance A to B

Write balanced equation and identify given and unknown quantities

- Given: number of moles of A you have

- Unknown: The substance B you want to convert to

Use molar ratios

- From the balanced equation, identify the molar ratios between A and B

- these ratios are determined by the coefficients in front of the substances

- if the balanced equation is  $a\text{A} + b\text{B} \rightarrow c\text{C} + d\text{D}$ , then the molar ratio of A to B is  $b/a$

Use the molar ratio to calculate the number of moles of substance B that will form or react with the moles of substance A

- Multiply the moles of A by the appropriate molar ratio

Ensure all reactants and products are included in the stoichiometric calculation:

- **Limiting and excess reactants:**

limiting: reactant completely consumed in reaction, limiting amount of product that can be formed

excess: reactant not completely consumed and is left over after limiting reactant is consumed fully

- **Theoretical yield**

Max. amount of product that can be obtained from given amount of reactants assuming all reactants are converted to products according to the stoichiometry of the balanced equation



By massiveballs

Not published yet.

Last updated 12th April, 2024.

Page 8 of 18.

Sponsored by **ApolloPad.com**

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>



### Moles of substance A to B (cont)

#### -compare actual and theoretical yields

actual yield: amount of product obtained from reaction in practice  
 if its less than the theoretical yield, it indicates one or more reactants were limiting and the reaction didnt proceed to completion.  
 if any reactant is limiting or in excess, adjust the stoichiometric calculations:  
 limiting: calculate amount of product formed based on its quantity  
 excess: determine amount left over after reaction is complete  
 To conclude, indicate the number of B moles formed or reacted based n the given A moles

### grams to atoms

find molar mass and calculate number of moles by dividing number of grams by molar mass  
 multiply the number of moles by avogadro's number  
 Example:  
 10 grams of Na to atoms  
 molar mass of Na is approx. 22.99 g/mol  
 divide 10 grams by molar mass  
 $= 10/22.99$   
 $= 0.435$  moles  
 Multiply number of moles by Avogadro's number  
 $0.434 \times (6.022 \times 10^{23})$   
 $= 2.6 \times 10^{23}$  atoms in 10 grams of Na

### Naming ionic main metal compounds

Some ionic compounds containing a transition metal require Roman numerals while some don't. The way we determine this is whether or not that transition metal can exhibit multiple oxidation states.  
 Elements in groups 1, 2 and 13 commonly have 1 oxidation state:  
 - Sodium, Na, is usually  $\text{Na}^+$  with a +1 charge, calcium, Ca, is usually  $\text{Ca}^{2+}$   
 - They don't require the use of roman numerals  
 For these kinds of ionic compounds, we simply use the same naming system as the other ionic compounds.

### Ionic compounds chemical formula

Determine if the elements in the compound typically form ions and which charge it forms.  
 Then, balance the charges  
 - since ionic compounds are electrically negative, the total positive charge from the cations must balance the total negative charge from the anions. To achieve this, you may need to adjust the number of ions present in the compound.  
 when writing the formula, simplify the subscripts by dividing them by their greatest common number, but don't change the ratio between them.  
 if there is more than one possible ionization state, use Roman numerals.  
 Example:  
 Sodium chloride ( $\text{NaCl}$ )  
 - Sodium, Na, has a +1 charge =  $\text{Na}^+$   
 - Chlorine, Cl, has a -1 charge =  $\text{Cl}^-$   
 - Since they have equal opposite charges, no need for adjustment/balancing  
 Aluminium chloride  
 - Al has a charge of  $+3$ , while chlorine has a charge of  $-1$   
 - So, we just need to swap the subscripts with the number of the other element present:  
 $\text{Al}^{+3} + \text{Cl}^{-1} \rightarrow \text{Al}_1\text{Cl}_3$  (we can omit the 1 from Al coz its already a given)  
 - Now, the total Al charge is +3, and the total Cl charge is -3, making them balanced  
 Aluminum chloride =  $\text{AlCl}_3$

### Grams to molecules

Find total molar mass of all elements in substance.  
 Use the conversion factor Avogadro's number -  $6.022 \times 10^{23}$  molecules/mol  
 Calculate number of moles  
 - divide number of grams by molar mass  
 Convert moles to molecules by multiplying the number of moles by Avogadro's number  
**10 g H<sub>2</sub>O to molecules**  
 Molar mass = 18.016g/mol  
 Divide 10 grams by molar mass  
 $= 10 \text{ grams} / 18.016 \text{ g/mol} = 0.555$   
 Convert to molecules  
 $0.555 \times (6.022 \times 10^{23})$



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 9 of 18.

Sponsored by **ApolloPad.com**

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Grams to molecules (cont)

=  $3.34 \times 10^{23}$  molecules of H<sub>2</sub>O

### Grams to moles

Find the atomic mass of each element and multiply it by the number of that element in 1 molecule of the substance.

Add the masses of all the atoms to find the molar mass.

Multiply the number of grams by the reciprocal molar mass of the substance.

Example:

50 grams of H<sub>2</sub>O to moles

- Molar mass of H<sub>2</sub>O is sum of the masses of H and O

- (H)  $1.008\text{g/mol} \times 2$  + (O)  $16\text{g/mol}$

= 18.016

Determine the substance: We want to convert grams of water (H<sub>2</sub>O) to moles.

- The reciprocal of the molar mass of H<sub>2</sub>O is  $1/18.016\text{g/mol}$

- multiply 50 grams by  $1/18.016\text{g/mol}$

= 2.78 moles

### Find amount of mols in gs of substance

Find molar mass of substance and divide it by the amount of grams

Example:

Moles in 1 gram H<sub>2</sub>O

- molar mass =  $18.016\text{g/mol}$

- Divide by 1

-  $1/18.016 = 0.0555$

Moles in 2 grams H<sub>2</sub>O

- multiply reciprocal of H<sub>2</sub>O by 2

=  $2 \times (1/18.016)$

= 0.1109 moles in 2 grams of H<sub>2</sub>O

### Naming polyatomic ions

Polyatomic ions are ions that contain more than 1 atom, monatomic ions only contain 1 atom

NO<sub>3</sub><sup>-</sup>, NO<sub>4</sub><sup>-</sup> = polyatomic

N<sub>3</sub><sup>-</sup> = monatomic

Suffixes:

'ate' or 'ite' - typically polyatomic ion that contains 1 oxygen atom

'ide' - typically monoatomic ion that lacks oxygen

so, N<sub>3</sub><sup>-</sup> is **Nitride**

and NO<sub>3</sub><sup>-</sup> is **Nitrate**

Example:

Cl<sup>-</sup> Chloride (no oxygen - ide)

ClO<sup>-</sup> Hypochloride (1 more oxygen - 'hypo' prefix)

ClO<sub>2</sub><sup>-</sup> Chlorite (1 more oxygen - ide)

ClO<sub>3</sub><sup>-</sup> Chlorate (1 more oxygen - ate)

ClO<sub>4</sub><sup>-</sup> Perchlorate (1 more oxygen - 'per' prefix)

### Dilution p2

**Water needed to dilute 100ml of 1.0 M solution of NaCl to a 0.25 solution**

Initial volume and concentration

100ml and 1.0 M respectively

Final concentration = 0.25 M (as desired)

Calculate amount of solute (initial)

Initial volume x initial concentration

=  $100\text{ml} \times 1.0\text{M} = 100$  moles

Calculate final volume needed

- we want to dilute solution to a final concentration of 0.25 M, so we use this formula

Final concentration = amount of solute (initial)/final volume

rearrange formula

Final volume = amount of solute (initial)/final concentration

=  $100 \text{ moles} / 0.25 = 400\text{ml}$

To get the amount of water needed, we calculate the difference btwn the final volume and the initial volume

Final volume - initial volume



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 10 of 18.

Sponsored by **ApolloPad.com**

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Dilution p2 (cont)

= 400ml - 100ml = 300ml

So you add 300ml of water

### Covalent/molecular compound chemical formula

#### Sulfur Dioxide

No prefix in front of 'sulfur' means there is only one in the compound

'di' prefix in front of 'oxygen' means there are 2 in the compound

So, the formula would be SO<sub>2</sub>

#### Dinitrogen pentoxide

'di' in front of nitrogen means there are two in the compound

'penta' in front of oxygen means there are 5 in the compound

So, the formula would be N<sub>2</sub>O<sub>5</sub>

### Grams to make solution

Number of moles can be calculated using the formula

$$n = c \times v$$

c - concentration in mol/L, V - volume in L

$$\text{mass} = n \times \text{molar mass}$$

How many grams of solid Mg(NO<sub>3</sub>)<sub>2</sub> are required to make 2.5 L of a 1.5 M Mg(NO<sub>3</sub>)<sub>2</sub> solution?

V = 2.5, the C = 1.5 mol/L, we want to find the mass of solid

Mg(NO<sub>3</sub>)<sub>2</sub> needed to make this solution

$$N = 1.5 \text{ mol/L} \times 2.5 \text{ L}$$

$$= 3.75 \text{ moles}$$

Now calculate the mass

$$3.75 \text{ moles} \times 148.31 \text{ g/mol}$$

$$= 556.1625 \text{ grams, rounded to 556 grams}$$

### Naming: ionic compounds

A compound is a substance with more than 1 element

NaCl is Sodium chloride - two different elements present, making it a compound

An ionic compound is one that is composed of ions

NaCl is composed of Na<sup>+</sup> and Cl<sup>-</sup>, an anion and a cation respectively

When naming an anionic compound, the ending of the last element is changed to 'ide'

### Naming: ionic compounds (cont)

NaCl - Sodium + chlorine = sodium chloride

Name the first element, and end the second element in 'ide'

Example:

AlP contains Aluminium and phosphorous = Aluminium phosphide

Polyatomic ionic compounds follow the same rules, look at the section for them

### Atoms in grams

Calculate molar mass of compounds then calculate the number of moles

$$n = \text{mass} / \text{molar mass}$$

multiply with avo number

sodium atoms in 1kg of Na<sub>2</sub>SO<sub>4</sub>

molar mass Na = 22.99

$$S = 32.07$$

$$O = 16$$

$$2 \times 22.99 + 32.07 + 4 \times 16$$

$$= 141.05 \text{ g/mol}$$

Moles of Na<sub>2</sub>SO<sub>4</sub>

$$= \text{mass} / \text{molar mass} = 1000 / 141.04$$

$$= 7.09$$

since there are 2 moles of Na for every 1 mole of Na<sub>2</sub>SO<sub>4</sub>, Na atoms are doubled

$$2 \times 7.09 = 14.18$$

$$14.18 \times 6.022 \times 10^{23}$$

$$\text{Na atoms} = 8.53 \times 10^{24}$$

### Amount of molecules in gs

Divide mass (g) by the molar mass of the molecule, then multiply it by avos number ( $6.022 \times 10^{23}$ )

Oxygen molecules in 6 grams of oxygen

$$n = \text{mass (g)} / \text{molar mass (g/mol}^1)$$

$$\text{molar mass of O}_2 = 16 \times 2 = 32$$

$$6 / 32 = 0.187$$

$$0.187 \times 6.022 \times 10^{23} \text{ molecules/mol}$$



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 11 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Product formed in reaction

Balance the equation, and determine the number of moles for each reactant

$N = \text{mass of reactant} / \text{molar mass}$

Barium peroxide reacts with hydrochloric acid to form peroxide, H<sub>2</sub>O<sub>2</sub>:

$\text{BaO}_2(\text{s}) + 2\text{HCl}(\text{aq}) \rightarrow \text{H}_2\text{O}_2(\text{aq}) + \text{BaCl}_2(\text{aq})$ .

If 2.01 g of barium peroxide is reacted with 0.75 g of acid (HCl) how much peroxide will be produced?

Determine moles for each reactant

BaO<sub>2</sub> barium peroxide

-  $N(\text{BaO}_2) = 2.01\text{g} / 169.3 \text{ g/mol}$

Hydrochloric acid HCl

-  $N(\text{HCl}) = 0.75\text{g} / 36.458 \text{ g/mol}$

Compare the number of moles

From the equation, 1/2 moles of BaO<sub>2</sub> react with 1 mole of HCl

$N(\text{BaO}_2) = 1/2 \times n(\text{HCl})$

Calculate the number of moles of BaO<sub>2</sub> using the same ratio

Compare the moles, whichever is smaller is the limiting reagent

HCl = 0.0199 moles

BaO<sub>2</sub> = 0.0103 moles

- HCl limiting reagent

Calculate product mass based on limiting reagent/HCl using stoichiometry to find number of moles

In the equation, the stoichiometric coefficients represent the mole ratio btwn reactants and products

HCl is limiting, and 2 moles of HCl react to produce 1 mole of H<sub>2</sub>O<sub>2</sub>

Therefore,  $n(\text{H}_2\text{O}_2) = 1/2 n(\text{HCl})$

Molar mass of H<sub>2</sub>O<sub>2</sub> is 34.014g/mol

To convert moles to grams, use its molar mass

- Number of moles x molar mass

- 0.0103 moles x 34.014g/mol

### Naming covalent/molecular prefixes

Mono - 1

Die - 2

Tri - 3

Tetra - 4

Penta - 5

Hexa - 6

Hepta - 7

Octo - 8

Nona - 9

Deca - 10

CO = Carbon monoxide

- 1st element has subscript of 1, but doesn't need prefix 'mono'

- 2nd element has subscript of 1 (1 oxygen atom), so prefix 'mono' is used

CO<sub>2</sub> = Carbon dioxide

- 2nd element has subscript of 2 (2 oxygen atoms), so 'di' is used

NO<sub>2</sub> = Nitrogen Dioxide

N<sub>2</sub>O<sub>5</sub> = Dinitrogen pentoxide

- 1st element has subscript of 2, so 'die' is used

- 2nd element has subscript of 5, so 'penta' is used

### molecular shape and molecule polarity

To predict molecular shape and whether a molecule is polar; identify central atom (which can form most bonds/least electronegative)

determine the electron geometry around the central atom by considering both bonding and non-bonding electron pairs

- NCl<sub>3</sub>, N has 1 lone pair and forms 3 single bonds with the Cl

- this gives it a tetrahedral electron geometry

Determine the molecular shape by considering only the positions of bonded atoms

- in NCl<sub>3</sub>, the lone pair of electrons on N repels the bonding pair, causing the molecule to adopt a trigonal pyramidal shape

Determine polarity - consider the electronegativity of the atoms and the molecular shape

- NCl<sub>3</sub>, N is less electronegative than Cl, meaning the bonds btwn N and Cl are polar



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 12 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### molecular shape and molecule polarity (cont)

- the chlorine atom carries a partial neg charge, and the N carries a partial pos charge

While the arrangement of atoms in  $\text{NCl}_3$  is symmetrical, symmetry alone doesn't determine polarity. The distribution of electron density due to the lone pair of N still results in a net dipole moment (overall polarity of the molecule), making the molecule polar.

### Moles produced and molecules to react



**If 5.23 moles of CO react with excess oxygen, how many moles of  $\text{CO}_2$  are produced?**

Moles of CO = 5.23 mol

need to find moles of  $\text{CO}_2$  produced

From equation, 2 moles of CO react with 1 mole of  $\text{O}_2$  to make 2 moles of  $\text{CO}_2$ , meaning ratio is 2:2  $\rightarrow$  1:1

Using the given moles of CO and the ratio, calculate moles of  $\text{CO}_2$  produced

$$\begin{aligned} \text{Moles of } \text{CO}_2 &= (\text{given moles of CO}) \times \frac{\text{moles of } \text{CO}_2}{\text{moles of CO}} \\ &= 5.23 \text{ mol} \times \frac{2 \text{ mol } \text{CO}_2}{2 \text{ mol CO}} \\ &= 5.23 \times 1 \end{aligned}$$

moles of  $\text{CO}_2$  = 5.23 mol

when 5.23 moles of CO react, 5.23 moles of  $\text{CO}_2$  are produced

**How many oxygen molecules were required to react all the  $\text{CO}$ ?**

The equation shows us that 2 moles of CO react with 1 mole of  $\text{O}_2$  to produce 2 moles of  $\text{CO}_2$ , so the ratio between CO and  $\text{O}_2$  is 2:1

$$\begin{aligned} \text{Moles of } \text{O}_2 &= (\text{given moles of CO}) \times \frac{\text{moles of } \text{O}_2}{\text{moles of CO}} \\ &= 5.23 \times \frac{1 \text{ mol } \text{O}_2}{2 \text{ mol CO}} \\ &= 5.23/2 \end{aligned}$$

= Moles of  $\text{O}_2$  = 2.615

Convert to molecules

1 mole of substance contains  $6.022 \times 10^{23}$  molecules, so we multiply 2.615 by this number

$$= 1.572 \times 10^{24}$$

exponent increases by 1 (23-24) bc when we convert moles to molecules, we are multiplying by Avogadro's number -  $6.022 \times 10^{23}$  molecules per mole.

if we have 2 moles of a substance, we have  $2 \times 6.022 \times 10^{23}$  molecules, which is  $1.2044 \times 10^{24}$  molecules

### Moles produced and molecules to react (cont)

- each additional mole increases the number of molecules in Avogadro's number

when we increase the number of moles by 1, the exponent of 10 increases by 1 in the scientific notation representation of the number of molecules

### Electronic configuration

Understand the subshells

Electronic configuration describes the distribution of electrons in the atomic orbitals of an atom

each subshell is labelled with the principal quantum number (n) and the orbital type (s,p,d,f)

1s<sup>2</sup>, 2s<sup>2</sup>, 2p<sup>4</sup>

- n is 1 for the 1s subshell, 2 is for the 2s and 2p subshell

- the value of n represents the energy level of the orbital. The superscript after each subshell represents the number of electrons in that subshell

- 1s<sup>2</sup>, 1s subshell is filled with 2 electrons

- 2s<sup>2</sup>, 2s subshell is filled with 2 electrons

- 2p<sup>4</sup>, 2p subshell filled with 4 electrons

To determine the element that corresponds to the electron configuration, you use the periodic table

find the element with the atomic number that matches the sum of the superscripts in the electron configuration

**Elements and ions in their ground state**

identify the atomic number (z) of an element

- carbon has a z of 6 - 6 protons and 6 electrons in its neutral state

determine the shell occupied by the valence electrons

- C has an atomic number of 6, so we fill the electrons into the available order of increasing energy

use the Aufbau principle

- electrons fill the lowest energy orbitals before moving to the higher energy ones

- fill the 1s, then the 2s, and then the 2p orbitals

For carbon (z=6), the electronic configuration is 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>2</sup>

- 2 electrons in 2s orbital, 2 electrons in the 2s orbital and 2 in the 2p orbital



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 13 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>

### Moles

1 mole =  $6.022 \times 10^{23}$   
 1 mol of carbon atoms =  $6.022 \times 10^{23}$  atoms of Carbon  
 1 mol of CO<sub>2</sub> =  $6.022 \times 10^{23}$   
 2 mol of carbon =  $2 \times 6.022 \times 10^{23}$   
 4 mol of C  
 4 mol C /  $1 \times 6 \times 10^{23}$  C atoms / 1 mol C  
 =  $4 \times 6 = 24$   
 =  $24 \times 10^{23}$   
 Move the decimals to the left <-- =  $10^{23}$  goes up by as many places as you moved to the left  
 Move decimals to the right --> it goes down

### Moles

The mass number of an element also represents the 'molar mass'  
 - 1 mol of an element has a mass of (its mass number)  
 - Nitrogen's mass number is 14 = 1 mol of N has a mass of 14g; 14g of N contains  $6.022 \times 10^{23}$  atoms  
 - Mole is proportional to its 'molar mass'  
 - 2 mol of N = 28g N  
 Figuring out the molar mass of a compound can be done by identifying the molar mass of each element present and adding them together  
**O<sub>3</sub>**  
 - 1 oxygen atom has a mass number of 16, so 6 of them would be  $3 \times 16 = 48$   
 - Molar mass of ozone/O<sub>3</sub> = 48g/mol  
**CO<sub>2</sub>**  
 - Carbon mass number is 12.01, Oxygen mass number is 16, so  $16 \times 2 = 32$   
 -  $32 + 12.01 = 44.01$ g/mol  
**Calcium phosphate**  
 - 3 calcium atoms, 2 phosphate groups with 1 P atom and 4 oxygen atoms  
 - first, balance the formula  
 - Calcium has 3 atoms  
 - 1 P atom in each phosphate group, since there are 2 groups, that's 2 P in total  
 - Each group has 4 O atoms, so  $4 \times 2 = 8$  O atoms in total

### Moles (cont)

= Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>  
 - Molar mass of C =  $40.08 \times 3 = 80.16$   
 - Molar mass of P =  $30.97 \times 2 = 61.94$   
 - Molar mass of O =  $16 \times 4 = 64$   
 Then you add them together  
 Molar mass of Calcium phosphate/Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> = 310.18g/mol

### oxidation state of an atom in a compound

#### General rules

- Oxidation state of an atom in its elemental form is always 0
- For monatomic ions, the oxidation state is equal to the charge of the ion
- The sum of the oxidation state of all atoms in a neutral compound is 0, and it equals the charge of the compound if it's an ion
- In compounds, some elements have fixed oxidation states (Group 1 metals always have an oxidation state of +1, group 2 metals always have an oxidation state of +2, oxygen is usually -2)

Start with the elements that have a fixed oxidation state/are in elemental form, and assign their states based on rules above  
 If an element has variable oxidation states, use these rules

- Assign the oxidation state of oxygen as -2, unless it's in a peroxide or when combined with fluorine, where it has a positive state
- Hydrogen usually has a +1 state, except when bonded with metals where it's -1
- Group 1 metals are +1, group 2 metals are +2, group 13 are +3 in compounds
- in compounds, fluorine is always -1
- the sum of oxidation states in neutral compounds is 0

After assigning states to each atom, check that the sum of the states is equal to the total charge of the compound or ion, and if it's neutral it should be 0

#### Example

NH<sub>3</sub>

H usually is +1, O is usually -2

The sum of the oxidation states should equal 0 as it's neutral  
 H is typically +1, so all 3 atoms in NH<sub>3</sub> will equal a +3 charge.



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 14 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>

### oxidation state of an atom in a compound (cont)

NH<sub>3</sub> is neutral, so the sum of oxidation states must equal 0, so Nitrogen must have a state that offsets the total charge from the H atoms - +3

N's state can be calculated by subtracting the sum of oxidation states of H from 0

Oxidation state of N is x

$$-x + 3(+1) = 0$$

$$-x + 3 = 0$$

$$-x = -3$$

Nitrogen's oxidation state therefore is -3

#### example

Nitrate ion has a -1 charge, and O has a -2 charge

lets say the nitrogen state is x

$$-3 \text{ O atoms, each } -2 - 3(-2) = -6$$

The sum of oxidation states must equal the charge of the ion, -1, so nitrogen must have an oxidation state that offsets the total charge of the oxygen atoms -6.

$$x + -6 = -1$$

$$x = -1 + 6$$

$$x = +5$$

so the oxidation state of NH<sub>3</sub> is -1, and the nitrate ion has a state of +5, while the O atom has a state of -2

### Hydrated ionic compound empirical formula

Identify the ionic compound and the number of water molecules in it Determine the masses of each component separately, including the mass of the anhydrous salt (w/o water) and the mass of the water molecules.

These can be find from the given total mass of the compound.

Calculate the molar mass

Determine molar mass of the anhydrous salt by summing the molar masses of each element in the compound

To find the molar mass of the anhydrous salt, sum each of the molar masses in the compound

- molar mass of compound = molar mass of each atom added together

Then, determine the molar mass of H<sub>2</sub>O

Calculate the moles

### Hydrated ionic compound empirical formula (cont)

Use the masses and the molar masses to find the number of moles in each component

For the anhydrous salt, use this formula:

- Moles of A.salt = mass of A.salt/molar mass of A. salt for the water:

- Moles of water = mass of water/molar mass of water

Determine the simplest ratio

divide the number of moles of each component by the smallest number of moles calculated, to get the simplest ratio of ions to water molecules

Round to whole numbers if not already whole numbers

Write the empirical formula using the whole number ratios

- the subscripts in the formula represent the number of ions or water molecules in one formula unit of the compound

#### Example

Copper (II) sulfate pentahydrate - CuSO<sub>4</sub> + 5H<sub>2</sub>O

- made of copper (II) sulfate - CuSO<sub>4</sub> - and 5 H<sub>2</sub>O/water molecules.

Lets say we have 250 grams, to determine the mass of the A.salt (CuSO<sub>4</sub>) and the mass of water by weighing the sample

Calculate molar masses

CuSO<sub>4</sub> molar mass = 159.55 g/mol

H<sub>2</sub>O molar mass = 18.02 x 5 = 90.10 g/mol

calculate moles

CuSO<sub>4</sub> = 100/159.55 = 0.627 mol

5 H<sub>2</sub>O = 50/90.10 = 0.554 mol

Determine the simplest ratio

divide the number of moles of each component by the smallest number of moles calculated

$$- 0.627/0.554$$

$$= 1.13$$

Round to nearest whole number

$$= 1$$

The ratio is therefore 1:5, so the empirical formula is CuSO<sub>4</sub> + 5H<sub>2</sub>O



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 15 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>

### Empirical formula from % composition

Convert % to grams

Start by assuming you have 100g sample of the compound, and convert the percentages of each element to grams

- if a compound contains 40% carbon, it means there are  $40/100 \times 100\text{g} = 40\text{g}$  of carbon in 100g of the compound

Convert grams to moles

use the molar mass of each element to convert to moles

- Moles = grams/molar mass

Determine simplest ratio

Divide number of moles of each element by the smallest number of moles calculated

- if the ratios obtained aren't whole numbers, then round them to the nearest whole number.

- if the numbers are close to whole numbers, you can multiply all ratios by the same number to make them whole

Write formula

use the whole number ratios to write the empirical formula of the compound.

the subscripts in the formula represent the number of atoms of each element in one molecule of the compounds

#### Example

Compound with 40% carbon, 6.7% hydrogen, and 53.3% oxygen by mass

Convert % to grams

C = 40.0g, H= 6.7g, O= 53.3g

Convert G to moles

Using the molar mass

- moles = mass (grams)/molar mass (grams/mol)

- C:  $40.0/12.01 = 3.33$  moles

- H:  $6.7/1.01 = 6.67$  moles

- O:  $53.3/16 = 3.33$  moles

Determine the simplest ratio

Divide the number of moles of each element by the smallest number of moles

The ratio of C:H:O = 1:2:1

They are close enough to whole numbers, and therefore don't need to be rounded.

### Empirical formula from % composition (cont)

The empirical formula would therefore be:

CH<sub>2</sub>O

#### Example

Analyses of a compound found it to contain, by mass: 63.68% C, 12.38% N, 9.80% H and 14.14% O. Calculate the empirical formula for this compound

Carbon:  $63.68/12.01 = 5.30$

N:  $12.38/14.01 = 0.884$

H:  $9.80/1.008 = 9.72$

O:  $14.14/16 = 0.900$

Divide number of moles each by the smallest number, which is 0.884 from Nitrogen

$5.30$  and  $8.884$  and  $9.72$  and  $0.900$  all divided by  $0.884$

= 6, 1, 11, 1 after simplifying

= empirical formula becomes C<sub>6</sub>NH<sub>11</sub>O<sub>1</sub>

Divide mass percentage of each element by its molar mass to find number of moles

Determine simplest whole-number ratio of moles by dividing each number of moles by the smallest number of moles

Write the empirical formula

Analyses of a compound found it to contain, by mass: 63.68% C, 12.38% N, 9.80% H and 14.14% O. Calculate the empirical formula for this compound

Carbon:  $63.68/12.01 = 5.30$

N:  $12.38/14.01 = 0.884$

H:  $9.80/1.008 = 9.72$

O:  $14.14/16 = 0.900$

Divide number of moles each by the smallest number, which is 0.884 from Nitrogen

$5.30$  and  $8.884$  and  $9.72$  and  $0.900$  all divided by  $0.884$

= 6, 1, 11, 1 after simplifying

= empirical formula becomes C<sub>6</sub>NH<sub>11</sub>O<sub>1</sub>



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 16 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>



### Mass of excess reactant

Write balanced equation, find limiting reactant, calculate theoretical yield of the product, and determine the excess reactant.

Now, calculate the amount left over by subtracting the amount of excess reactant that reacted from the initial amount of excess reactant given.

Convert to mass

Use the molar mass of the reactant to convert the amount of excess reactant left from moles to grams.

### Dilution

First determine initial volume and concentration of the solution before dilution

Then determine the final volume of the solution after dilution by adding the initial volume of the solution to the volume of the solvent added during dilution

Calculate the initial amount of solute by using the initial volume and concentration to calculate the amount of solute (substance being diluted) present in the solution before dilution.

- Amount of solute (initial) = initial volume x initial concentration

Now determine the final concentration

- Final concentration = amount of solute (initial)/final volume

**if we dilute 100ml of a 1.0 M (concentration) of KCl to 400ml with 300ml of water, what will the final concentration be?**

Initial volume = 100ml

initial concentration = 1.0 M

Final volume = initial volume + volume of solvent added

= 100ml + 400ml = 500ml

Use the initial volume and concentration to calculate the amount of solute (KCl) present in the solution before dilution

- initial solute amount = initial volume x initial concentration

= 100ml x 1.0 M = 100 moles

Final concentration

= amount of solute (initial)/final volume

= 100 moles/500ml = 0.2 M

### A grams - B grams

Balance equation and identify given and unknown qualities.

Convert grams of A to moles

use the given mass of A and it's molar mass

- moles = mass (grams)/molar mass (grams/mol)

Use the molar ratios

Convert moles of A to B by using the molar ratio

- multiply A moles by appropriate molar ratio

Convert moles of B to grams

- mass (grams) = moles x molar mass (grams/mol)

Conclude by showing the mass of B formed or reacted based on the given mass of A

### Stoichiometry: Moles Formed in Reaction

Write balanced chemical equation

Identify the given and unknown qualities

- given: initial amount of moles

- unknown: moles that will be formed

Use the molar ratio to determine the answer

Example:

How many moles of SO<sub>3</sub> will form when 3.4 moles of sulfur dioxide react with excess oxygen gas?

Write the balanced equation

Balance the equation for the reaction between SO<sub>2</sub> and O<sub>2</sub> to form sulfur trioxide

- SO<sub>2</sub> + O<sub>2</sub> → SO<sub>3</sub>

- 2SO<sub>2</sub> + O<sub>2</sub> → 2SO<sub>3</sub>

Given: 3.4 moles of SO<sub>2</sub>

Unknown: moles of SO<sub>3</sub> formed

Use the molar ratio:

The balanced equation shows us that 2 mole of SO<sub>2</sub> react to form 2 moles of SO<sub>3</sub>, which means the molar ratio between them is 1:1.

Therefore, if 3.4 moles of SO<sub>2</sub> react, 3.4 moles of SO<sub>3</sub> will form

The reaction proceeds to completion, meaning all of the reactant

(SO<sub>2</sub>) is consumed and converted into products - there no limiting

factors that would prevent the complete consumption of the reactant.



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 17 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>

### Stoichiometry: Moles Formed in Reaction (cont)

In the reaction btwn SO<sub>2</sub> and O<sub>2</sub> to form SO<sub>3</sub>, if SO<sub>2</sub> is provided in excess, it implies theres enough oxygen to completely react with the SO<sub>2</sub> presented. Excess oxygen ensures that all the SO<sub>2</sub> molecules will find oxygen molecules to react with, thus the reaction can occur till all the SO<sub>2</sub> is consumed.

Because the reaction proceeds to completion and all the SO<sub>2</sub> is consumed, the number of SO<sub>3</sub> moles formed will be equal to the number of SO<sub>2</sub> moles initially present.

Therefore, the number of moles of SO<sub>3</sub> formed will also be 3.4 moles.

- 3.4 moles of SO<sub>3</sub> will form when 3.4 moles of SO<sub>2</sub> react with excess O<sub>2</sub>.

### Dilution p3

#### Dilute 250 mL of a 0.100 M solution from a 2.00 M solution

Initial volume = ?

Initial concentration = 2.00

Final volume = volume of diluted solution (250ml)

Final concentration = 0.100m

Calculate amount of initial volume

- since its unknown, we cant directly calculate the amount of solute from it, but we know that the amount of solute in the stock solution is equal to the amount of solute in the diluted solution after dilution (since no solute is added or removed during dilution)

So we can calculate the amount of solute using the final concentration and volume of the diluted solution.

Amount of initial solute = final concentration x final volume

= 0.100 M x 250mL = 25 moles

To find out how much of the stock solution (2.00M) we need to dilute to obtain the desired amount of solute (25 moles), we use this formula

- Final volume = Amount of solute (initial) / Initial concentration  
= 25 moles / 2.00 M = 12.5 mL

Calculate amount of water needed

this is the difference btwn final volume of the stock solution and the volume of the diluted solution

Water needed = Final volume of the stock solution - Final volume of the diluted solution

= 12.5 mL - 250 mL = -237.5 mL

### Dilution p3 (cont)

The amount of water is negative so we dont need to add additional water to the stock solution for the desired concentration of 0.100M  
We just remove 12.5mL of the stock solution and then add water to make up the remaining volume to reach 250mL, giving us the desired diluted solution with a concentration of 0.100M

Formula for dilution is  $C_1V_1 = C_2V_2$ , C<sub>1</sub> and V<sub>2</sub> are initial concentration and volume, and C<sub>2</sub> V<sub>2</sub> are final concentration and volume

Rearrange the formula to solve for v<sub>1</sub>

$V_1 = C_2V_2/C_1$

A laboratory technician is required to prepare 1.00 m<sup>3</sup> of 0.100 M H<sub>2</sub>SO<sub>4</sub>. What volume of 10 M H<sub>2</sub>SO<sub>4</sub> is required?

C<sub>1</sub> = 10M, V<sub>2</sub>=1.00m<sup>3</sup>, C<sub>2</sub> = 0.100M

$V_1 = 0.100M \times 1.00m^3 / 10M$

= 10L



By massiveballs

[cheatography.com/massiveballs/](https://cheatography.com/massiveballs/)

Not published yet.

Last updated 12th April, 2024.

Page 18 of 18.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>