

Refeeding Syndrome Cheat Sheet by Michellephillipso2 via cheatography.com/214485/cs/46707/

What is it?

May occur when a pt begins eating an adequate amount – oral, EN or PN - after a period of prelonged starvation or malnutrition

- Starvation Phase: ↓ insulin secretion, ↑
 fat and protein metabolism, ↓ Phosphate,
 K, Mg stores depleted (serum levels may be normal)
- Refeeding Phase: Sudden increase in carbohydrate intake → ↑ insulin secretion.
 Rapid intracellular shift causes ↓ serum electrolyte levels.

At risk

BMI <18.5, 10% weight loss 3-6/12, little/no intake >5 days, ↓ K, Mg, PO

chronic	chronic alcoholism/subs-
malnutrition	tance abuse
EDs	oncology
post-op	severe mental health

disorders

S/S	
Water/salt retention (oedema)	Impaired muscle contraction,
tachy/brady	glucose intolerance
respiratory difficulties	seizures
confusion	coma
hlurred vision	

By Michellephillips02

cheatography.com/michellephillips02/

Biochem

Consequences

Electrolyte imbalances (hypophosphataemia, hypokalaemia, hypomagnesaemia)

Abnormal glucose metabolism - hypoglycaemia or hyperglycaemia

Thiamine deficiency

Cardiac: arrhythmias, heart failure

Respiratory: muscle weakness, respiratory

Neurologic: confusion, seizures

Haematologic: anaemia, impaired immune

function

Intervention

Initiate feeding once: 1. RFS Supps (PO4, K, Mg, B1) have commenced. 2. Electrolytes are monitored. 3. Abnormalities are corrected

High risk pts: Start rate = 50% of goal *OR* Commence low-CHO feed @ ~6000kJ/day. Increase by 2000kJ/day until goal is met.

Lower risk pts: Start @ 1800kcal/day. Gradually increase when biochem is stable, eg +400cal/week

Avoid excessive Na (water retention)

NGT - BMI < 14kg/m2

Not published yet.

Page 1 of 1.

Last updated 11th July, 2025.

Sup 500mg PO4 bd. 100mg thiamine od for first week.

Guidelines & References

Handbook p169

ASPEN consensus recommendations for refeeding syndrome (2020)

ESPEN basics in clinical nutrition: Refeeding syndrome (2010)

Monitoring

Monitor weight daily

Electrolytes 6-8 hours after first re-feeding

Daily bloods for the first week

Second week = bloods 3/week

BGL 2/day

Sponsored by **CrosswordCheats.com**Learn to solve cryptic crosswords!

http://crosswordcheats.com