

### Specimens in the Lab

#### Adequate for PCR

- |                     |                               |
|---------------------|-------------------------------|
| - Whole Blood       | - Dried Blood                 |
| - Bone Marrow       | - Saliva                      |
| - Tissue/ Scrapings | - Bone, Teeth                 |
| - Bacteria          | - Amniotic fluid              |
| - Fungi/Yeasts      | - Hair follicles/ Hair Shafts |
| - Buccal Cells      | - Buccal Cells                |
| - Hair              | - Cerebrospinal fluid         |
| - Urine             | - Fixed tissue                |
| - Stool             | - Stool                       |
|                     | - Soil                        |
|                     | - Urine                       |

### DNA Extraction Methods

#### Organic

Uses organic chemical, phenol, chloroform

- |                                    |                                      |
|------------------------------------|--------------------------------------|
| 1. Lysis (NaOH, SDS)               | 2. Acidification (acetic acid, salt) |
| 3. Extraction (phenol, chloroform) | 4. DNA precipitation (ethanol)       |

#### Inorganic

Uses inorganic chemicals, detergent, ethylenediamine tetraacetic acid (EDTA), acetic acid, salt (salting out, spooling)

- |                                    |   |
|------------------------------------|---|
| 1. Lysis (Tris, EDTA, SDS)         | 2. Protein precipitation (sodium acetate) |
| 3. DNA precipitation (isopropanol) |   |

#### Solid phase

DNA is immobilized on a solid support, beads, or columns

- |                              |                                      |
|------------------------------|--------------------------------------|
| 1. Lysis (supplied reagents) | 2. Acidification (supplied reagents) |
| 3. Adsorption (low pH)       | 4. Wash (supplied buffer)            |
| 5. Elute DNA (low salt)      |                                      |

### DNA Extraction Methods (cont)

- |                       |                        |
|-----------------------|------------------------|
| Solid-Phase Isolation | -Kits<br>-Spin Columns |
|-----------------------|------------------------|

Nucleic acid binding to silica beads is the basis for many automated extraction systems

Limiting specimens (fixed tissue, dried, bone)

Rapid extraction for routine testing

#### Crude lysis

DNA does NOT need to be pure RE digestion, gel electrophoresis, screening large amounts of DNA, some PCR, challenging specimens

Proteolytic enzymes Proteinase K (PK)

Chelating resins Chelex

#### Chelex resin

Heat tissue (hair roots, saliva, etc.) in 300 uL 5%-20% Chelex 100 resin

Chelex removes multivalent cations

DNA is in supernatant

#### Fixed tissue

- |                                       |                              |
|---------------------------------------|------------------------------|
| 1. ~20 micron sections                | Macrodissect or microdissect |
| 2. Digest (proteinase K, Tris buffer) |                              |

### DNA Isolation

DNA isolation, purification and quantitation	Free of proteins, lipids, other nucleic acids
--	---

Different methods depending on specimen type	??
--	--

Different methods depending on type of DNA needed	??
---	--



### DNA Quantitation

#### Spectrophotometry

Instrument that allows you to measure the amount of light that is absorbed in a sample or the amount of light transmitted based on the solute in the sample

Filters and/or prisms are used to filter the light to obtain a certain wavelength

A= absorbance

%T= transmittance

#### A260/A280 Ratio

$1.65 \leq A260/A280 \leq 2.50$

dsDNA concentration =  $50 \text{ ug/mL} \times OD260 \times \text{dilution factor}$

#### Beer - Lambert Law

Absorbance is directly proportionate to the concentration of the solute (DNA)

Absorptivity constant - 50 for DNA

- Conversion factor from optical density unit (absorbance units) to concentration

- Residual phenol - bad!!

#### Calculations

Usually have to dilute samples from preps to begin

A DNA prep diluted 1:100 has an A260 reading of 0.200. Find the concentration in ug/mL

A DNA prep diluted 1:50 has an A260 reading of 0.307. Find the concentration in ug/mL.

### Separating out the cells

Differential Density gradient centrifugation

### Breaking open the cells

Alkaline lysis

Differential lysis

### Cell wall digestion

Lysozyme

Zymolyase

Detergents

Boiling



By **Morghay123**

[cheatography.com/morghay123/](https://cheatography.com/morghay123/)

Not published yet.

Last updated 8th September, 2018.

Page 2 of 2.

Sponsored by **ApolloPad.com**

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>