

### ACS (Acute Coronary Syndrome)

#### Definition

Spectrum of problems ranging from unstable angina to MI

#### Symptoms

Crushing pain/pressure; radiation to jaw, back, and left arm; SOB, diaphoresis, N/V; impending sense of doom

#### Most common etiology of MI

Preexisting atherosclerotic plaque → thrombus formation → prolonged myocardial ischemia → MI

#### EKG changes

Acute MI: progression from peaked T- waves → ST-deg ment elevation/depression → Q-wave → T-wave inversions (hours-days)

#### Laboratory Tests

EKG (within 10 min), troponin levels, CK:CK-MB ratio, MRI with gadolinium

#### Stable Angina

Reproducible pain, improved with rest, lasts <10min,

#### UA/NSTEMI

Occurs more often with less activity, not relieved with NTG, lasts >10min, EKG changes

#### STEMI

Complete occlusion, EKG changes

### ACS (Acute Coronary Syndrome) (cont)

#### Initial Treatment

ONAM: Oxygen, +/- nitro, antiplatelets (ASA+P2Y12), morphine (PRN), EKG monitoring, IV access

#### Discharge Treatment

- 1) ASA (life)
- 2) P2Y12 (1yr)
- 3) Statin
- 4) β-blocker
- 5) ACE-I
- 6) Aldosterone antagonist

#### Emergency Intervention

Door-to-needle within 30min; door-to-balloon within 90min

### Congestive Heart Failure (CHF)

#### Definition

Inability of the heart to keep up the the demands on it and pump blood with normal efficiency

#### Result of one or more of the following

Contractile ability of heart muscle, preload and after load of the ventricle, and heart rate

#### Etiology

MI, pericardial disorders, valvular disorders, congenital abnormalities, and non cardiac causes (high-output heart failure from thyrotoxicosis or severe anemia)

#### Clinical features of HFpEF/LHF

Exertional dyspnea, non-productive cough, fatigue, orthopnea, PND, basilar rales, gallops, exercise intolerance

### Congestive Heart Failure (CHF) (cont)

#### Clinical features of HFpEF/RHF

Distended neck veins, hepatic congestion, nausea, dependent pitting edema, \*edema + hepatomegaly, (R-sided failure often caused by L-sided failure)

#### Treatment

- 1) Loop
- 2) ACE-I
- 3) β-blocker
- 4) Spironolactone
- 5) Hydralazine + ISDN (esp in blacks)

### Hypertension

#### Primary HTN

Causes 95% of cases of HTN; multifactorial pathogenesis (genetics, salt, obesity, RAAS, NSAIDs, smoking, lack of exercise, metabolic syndrome)

#### Secondary HTN

Narrowing of aorta, RAS, chronic steroids, Cushings syndrome, pregnancy, thyroid and parathyroid disease, primary hyperaldosteronism, parenchymal renal dz)

#### Treatment Goal

All ages with DM or CKD ≤140/90  
Ages <60yo ≤140/90  
Ages ≥60yo ≤150/90

#### Treatment

First line: ACE-I/ARB, CCB, thiazides  
Other: alpha blockers, clonidine, guanfacine, hydralazine, minoxidil,

### Ischemic Heart Disease

#### Definition

Characterized by insufficient oxygen supply to cardiac muscle

#### Etiology

1) Atherosclerotic narrowing (most common). 2) Constriction of coronary arteries. 3) (Rare) congenital, emboli, arteritis, dissection

#### Risk Factors

Metabolic syndrome, male, older age, smoking, FmHx, HTN, DM, low-estrogen state, abdominal obesity, inactivity, dyslipidemia, EtOH, low fruits/veggies (cocaine → MI)

#### Un/stable Angina

See Above

#### Prinzmetal's (Variant) Angina

Caused by vasospasm at rest, exercise capacity preserved. Treated with CCBs, avoid β-blockers

#### EKG Findings

Horizontal or downsloping ST-segment depression

#### Treatment

Lifestyle changes, nitrates (nitro and LA), B-blockers, CCB, Ranolazine, ASA/Clopidigrel, revascularization