

Phases of the cell cycle Mitosis

Prophase	Chromosomes condense and spindle apparatus forms
Prometaphase	Kinetochores assembled at centromere, 2 opposite sides connected to microtubules
Metaphase	Lined up on imaginary metaphase plate. Polar microtubules extend from each spindle, overlap in middle, pole-pole connection
Anaphase	Cohesions are cleaved, daughters to opposite sides of cell. poles pulled apart
Telophase	Nuclear envelope reforms, chromosomes begin to condense
Cytokinesis	Division of cytoplasm

Types of dominance

Incomplete dominance	Phenotypes are blended together ex. pink flowers come from red and white alleles
Co-dominance	Both phenotypes show up ex. polka dots

Mitosis

Mitosis	When cells divide, two genetically identical sister cells are their products
Uses	Somatic cells

G1 checkpoint

1. Cells big enough
2. Sufficient nutrients
3. social signals present
4. Cells undamaged

G2 checkpoint

1. No errors in replication
2. Activated MPF (cyclin + CDK) present
3. Undamaged

Metaphase checkpoint

1. Chromosomes attach to spindles
2. Chromosomes properly segregated
3. MPF absent

Mechanisms of cell cycle progression

Nucleotide excision repair	1. Error detected in DNA by proteins 2. DNA nicking (cut at both sides of damage) 3. Helicase unwinds and removes region with damaged bases 4. DNA polymerase fills gap with undamaged strand as template 5. Nucleotide linkage (DNA ligase links the strand into existing strand. If successful continues past G1 checkpoint
P53 gene	Creates CDK inhibitors if the cell is damaged so if cyclin is still present, CDK can still say no if damaged
UVRA	recognizes DNA damage, signals to start repair, if damage can't be repaired cell won't divide anymore.
recA	Facilitates DNA repair



By NoelleEvelyn

cheatography.com/noelleevelyn/

Not published yet.

Last updated 14th March, 2025.

Page 1 of 3.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!

<https://apollopad.com>

Genes on X-chromosome

The X chromosome is larger	it holds most all of the sex-linked traits
In females	Females have 2 copies of X chromosome
	When sex linked traits are recessive they would need 2 copies to express the mutation
In males	Males only have one X-chromosome
	Males only need 1 copy of recessive X-linked trait to express the mutation

Segregation and Independent assortment

Law of segregation	Each diploid parent forms a haploid gamete
Independent assortment	Allels of different genes separate independently of each other to form gametes

Epistasis

Epistasis	The expression of one gene influences or masks the expression of another gene
Ex.	Fur color in golden retrievers

Map distance for F2 generation

Greater than 50 map units	Independently assorting
---------------------------	-------------------------

Independent assortment

Linked genes	Do not follow rules of independent assortment
	Too close together on chromosome to separate
Closer genes are	More likely they are linked
Independent assortment	occurs between chromosomes not within

Reciprocal vs Test cross

Reciprocal cross	The cross between a male with one phenotype and a female with another and then flipping
	Determines if sex plays a role in inheritance
Test Cross	Dominant phenotype crossed with recessive genotype
	Determines genotype of dominant phenotype

Genes arranged on chromosomes within genome

Karyotype	# and visual appearance of gametes
Genes hold	Instructions for making mRNA
Homologous chromosomes	Same genes in same location, but different versions of gene
Allels	versions of genes
Genotype	Allels present
Gene locus	location of genes

Asexual vs Sexual reproduction

Asexual	Sexual
No variation, exact clones	More variation
Quicker	Slower
Binary fission	Humans
Mitosis	Meiosis

Importance of Telomeres

Protect from	important DNA being cut out
Everytime cell divides	become shorter
Replication limit	prevents cancer
Why?	There is no 3' hydroxyl at end of lagging strand.
What?	G-rich series of repeats
Telomerase	elongates parental in 3' to 5' direction.



By NoelleEvelyn

Not published yet.
Last updated 14th March, 2025.
Page 2 of 3.

Sponsored by **ApolloPad.com**
Everyone has a novel in them. Finish Yours!

Both leading and Lagging strands

Single stranded binding proteins (SSBs)	Keep strands from attaching back together
Ligase	Fills in gaps or breaks in phosphodiester bonds of backbone
Helicase	Separates, unwinds double stranded DNA
Topoisomerase	Helps with stress on wound DNA, ex. Gyrase

DNA synthesis in lagging strand

Synthesized	in fragments (Okazaki fragments)
Initiated by	RNA polymerase
RNA polymerase	builds primers
DNA polymerase	replicates DNA off of primers
RNA primer	popped out of gaps and replaced with DNA polymerase

DNA synthesis in Leading strand

Synthesized	Continuously
Begins with	RNA primer
After RNA primer	DNA polymerase



By NoelleEvelyn

cheatography.com/noelleevelyn/

Not published yet.

Last updated 14th March, 2025.

Page 3 of 3.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>