

Cheatography

Algebra, Calculus I & II Cheat Sheet

by pablocdch via cheatography.com/30131/cs/8927/

Derivatives

Expression	Derivative
$f(x)=k$	$f'(x)=0$
$f(x)=x$	$f'(x)=1$
$f(x)=x^a$	$f'(x)=ax^{a-1}$
$f(x)=a^x$	$f'(x)=a^x \ln a$
$f(x)=kx$	$f'(x)=k$
$f(x)=1/x$	$f'(x)=-1/x^2$
$f(x)=\ln x$	$f'(x)=1/x$
$f(x)=e^u$	$f'(x)=u'.e^u$
$f(x)=\log_a x$	$f'(x)=1/x \ln a$
$f(x)=\sin(x)$	$f'(x)=\cos(x)$
$f(x)=\sec(x)$	$f'(x)=\sec(x).\tan(x)$
$f(x)=\arcsen(x)$	$f'(x)=1/\sqrt{1-x^2}$
$f(x)=\cos(x)$	$f'(x)=-\sin(x)$
$f(x)=\csc(x)$	$f'(x)=-\csc(x).\cot(x)$
$f(x)=\arccos(x)$	$f'(x)=-1/\sqrt{1-x^2}$
$f(x)=\tan(x)$	$f'(x)=\sec^2(x)$
$f(x)=\cot(x)$	$f'(x)=-\csc^2(x)$
$f(x)=\arctan(x)$	$f'(x)=1/1+x^2$

Integrals

$\int x^k dx$	$(x^{k+1})/(k+1) + C$
$\int x^{-1} dx$	$\ln x + C$
$\int dx$	$x + C$
$\int kF(x)dx$	$k \int F(x)dx$
$\int [F(x) \pm G(x)] dx$	$\int F(x)dx \pm \int G(x)dx$
$\int e^{kx} dx$	$(1/k)e^{kx} + C$
$\int a^{kx} dx$	$a^{kx}/[k \ln(a)] + C$
$\int \sin(kx) dx$	$-(1/k) \cos(kx) + C$
$\int \cos(kx) dx$	$1/k \sin(kx) + C$
$\int \sec(kx) dx$	$1/k (\ln \sec(kx) + \tan(kx)) + C$
$\int \tan(kx) dx$	$-(1/k) \ln \cos(kx) + C$
$\int \csc(kx) dx$	$1/k \ln \csc(kx) - \cot(kx) + C$
$\int \cot(kx) dx$	$1/k \ln \sin(kx) + C$
$\int \sec(kx) dx$	$1/k \sec(kx) + C$
$\int \csc(kx) \cot(kx) dx$	$-1/k \csc(kx) + C$
$\int \sec^2(kx) dx$	$1/k \tan(kx) + C$
$\int \csc^2(kx) dx$	$-1/k \cot(kx) + C$

Quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Factoring

$(a+b)^2$	$a^2 + 2ab + b^2$
$(a-b)^2$	$a^2 - 2ab + b^2$
$a^2 - b^2$	$(a-b)(a+b)$
$(a+b)^3$	$a^3 + 3a^2b + 3ab^2 + b^3$
$(a-b)^3$	$a^3 - 3a^2b + 3ab^2 - b^3$
$a^3 - b^3$	$(a-b)^3 + 3ab(a-b)$
$a^3 + b^3$	$(a+b)^3 - 3ab(a+b)$

Factoring

Given $x^2+ax+b=0$, then you have to find two numbers that when **multiplied** give you **B** and **added** give you **A**.

Example:

x^2+4x+3 , turns into: $(x+3)(x+1)$

Absolute value properties

$ x >a$	$x>a$ or $a<-x$
$ x <a$	$-a < x < a$

Divisions with 0

0/n	0
n/∞	0
n/0	∞

Logs properties

$\log_a B^n$	$n \log_a B$
$\log_a A = \ln e$	1
$\log_a 1 = \ln 1$	0
$\log_a(m \cdot n)$	$\log_a m + \log_a n$
$\log_a(m/n)$	$\log_a m - \log_a n$

Exponential properties

Trigonometrical identities

$\sin^2 \alpha + \cos^2 \alpha = 1$	
$\sec^2 \alpha = 1 + \tan^2 \alpha$	
$\cosec^2 \alpha = 1 + \cot^2 \alpha$	
$\tan \alpha = \sin \alpha / \cos \alpha$	
$\tan^2 \alpha + 1 = \sec^2 \alpha$	
$\cot^2 \alpha + 1 = \csc^2 \alpha$	
$\sin(x+y) = \sin(x) \cos(y) + \cos(x) \sin(y)$	
$\cos(x+y) = \cos(x) \cos(y) - \sin(x) \sin(y)$	
$\tan(x+y) = (\tan(x) + \tan(y)) / (1 - \tan(x) \tan(y))$	
$\sin(x-y) = \sin(x) \cos(y) - \cos(x) \sin(y)$	
$\cos(x-y) = \cos(x) \cos(y) + \sin(x) \sin(y)$	
$\tan(x-y) = (\tan(x) - \tan(y)) / (1 + \tan(x) \tan(y))$	
$\sin(2x) = 2 \sin(x) \cos(x)$	
$\cos(2x) = \cos^2(x) - \sin^2(x)$	
$\cos(2x) = 2 \cos^2(x) - 1$	
$\tan(2x) = (2 \tan(x)) / (1 - \tan^2(x))$	
$\sin^2(x) = 1/2 (1 - \cos(2x))$	
$\cos^2(x) = 1/2 (1 + \cos(2x))$	
$\sin(x) \cos(x) = 1/2 \sin(2x)$	
$\sin(x) \sin(y) = 1/2 (\cos(x-y) - \cos(x+y))$	
$\sin(x) \cos(y) = 1/2 (\sin(x-y) + \sin(x+y))$	
$\cos(x) \cos(y) = 1/2 (\cos(x-y) + \cos(x+y))$	
$\csc(x) = 1/\sin(x)$	
$\sec(x) = 1/\cos(x)$	
$\cot(x) = \cos(x)/\sin(x)$	
$\sin(-x) = -\sin(x)$	
$\cos(-x) = \cos(x)$	
$\tan(-x) = -\tan(x)$	

a^0	1
a^1	a
$a^m \cdot a^n$	a^{m+n}
a^m / a^n	a^{m-n}
$(a.b)^n$	$a^n \cdot b^n$
$(a/b)^n$	a^n/b^n
$(a^m)^n$	$a^{m.n}$
$a^{n/m}$	raiz m de a^n
a^{-1}	$1/a$

C

By **pablocdch**

cheatography.com/pablocdch/

Not published yet.

Last updated 9th January, 2017.

Page 1 of 2.

Sponsored by **ApolloPad.com**

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>