
Isotopes

Isotope - An atom with the same number of protons but a different number of neutrons.
Radioisotope - An isotope that is radioactive and sometimes unstable. They
decay.

Formulas

W1= q ΔV1
(J)..(.c)..(v)
Power - rate of doing work
P= w/t (joules/seconds)
Power w/t = qΔv/t
P=iΔv
Unit of energy
w=pt
(joules) = Watt*Sec
New energy unit = kWh
Electrical Energy
1kWh ≡ 1000*3600
1kWh ≡ 3.6*10^6J

resistance



1/Rt = 1/R1 + 1/R2 + ...+1/Rn if in parallel.

Electrical energy and power

Electrical energy (Joules) = potential drop (volts) x current (amps) x times (seconds) E=VIt
P=E/t were one watt = 1 joule per second
E/t = VIT/t or P= VI
Power (watts) = voltage (volts) x current (amps)
P=VI
How much energy does a 100W light bulb use in half an hour?
P=100W and t=0.5h
So E=100W x 0.5h = 50Wh or 0.05kWh
To find power used Volts times Amps

Resistance

$$R = \frac{pL}{A}$$

p = *resistivity of a material*
L = *length of the wire*
A = *cross sectional area of wire*


Ohm's law: ΔV=iR or V=Amps*Ohms

A charge q moving through a potential difference ΔV will lose potential energy: ΔU=qV

Electric field


The electric field in any region of space is defined as the electric force per unit charge: E = F/q
the force on a charge of q in an electric field is given by F= qE

Neutron Bombardment



Sponsored by **ApolloPad.com**
Everyone has a novel in them. Finish Yours!
https://apollopad.com

Decay



Radiaiton

Alpha radiation - Helium nucleus (2 protons and 2 neutrons), highly ionising, charge of 2+, heavy.

Beta radiation - fast moving electron, negatively charged electron, not as high ionising capabilities, charge of -1.

Gamma radiation - electromagnetic radiation, less ionising then alpha or beta.

Detecting radiation - Geiger-Muller tube counts ions that are produced inside it, these create an electrical signal.

Radiation in the body - Ionising radiation can damage or kill the cells, dna can replicate in the damaged form, cancers can form.

Ionising Radiation

Ionising means that the radiation rips off electrons from nearby atoms that it passes.

Three types of radiation: Alpha, Beta and Gamma


Alpha passes through paper.

Beta passes through thin metal.

Gamma passes through thick metal.

Alpha radiation is most ionising, so it loses it's energy very quickly.

Atomic Notation



Half life

Half-life - the time taken for the radioactivity of a specified isotope to fall to half its original value.

Nuclear fission/fusion

Fission- When a nucleus splits into two or more pieces usually after bombardment by neutrons.

Fusion- A process taking place inside stars in which small nuclei are forced together to make larger nuclei. Energy is released in the process.

Chain reaction - A series of nuclear fissions that may or may not be controlled. The neutrons that are released cause the reaction.

Nuclear Fission Reactors

Used to harness energy from Fission reactions.

Neutrons released from Uranium-235 when it undergoes fission are travelling at high speeds, this leads to a chain reaction which causes an explosion.

The heat generated from the fission process is used to make steam which drives the turbine.


Fuel rods- long, thin rods containing pellets of enriched uranium

moderator- material that slows neutrons.

control rods- rods made of a material that absorbs neutrons

coolant- a liquid or gas to absorb the heat energy

Nuclear Reactor



Electric charge

Conductors: All metals, especially silver, gold, copper aluminium and any ionic solution.

Moderate conductors: Water and earth.

Semi-conductors: Silicon, Germanium and skin.

Insulators: Plastics, polystyrene, dry air, glass, porcelain, cloth (dry)

Moderate insulators: wood, paper, damp air, ice and snow.

Sponsored by **ApolloPad.com**
Everyone has a novel in them. Finish Yours!
https://apollopad.com

Electrical forces and fields

$$F = \frac{kq_1q_2}{r^2}$$

For the forces between two charges q1 and q2 at a distance of r
k= 9.0 x 10^9 N m^2 x C^2

Electric Current

Electric current is the rate of transfer of charge: I=q/t
where q is the charge transferred and t is the time taken.
1 ampere (A) = 1 coulomb per second ($C*s^{-1}$)
So 1 coulomb (C) = 1 ampere second (A*s)
1 volt = 1 joule per coulomb ($1V = 1JC^{-1}$)
1 ohm = 1 volt per ampere ($1ohm = 1VA^{-1}$)


Resistance

ΔVBattery=i Rtotal
ΔvBattery=ΔV1+ΔV2
Therefore i*Rtotal = ΔV1+ ΔV2=iR1+iR2
Rtotal = R1 + R2
R= V/I or V = IR

Electric Circuits

In any electric circuit the sum of all currents flowing into any point is equal to the sum flowing out of it.
The total potential drop around a closed circuit must be equal to the total EMF (electromotive force, the energy provided by the cell)

Symbols and devices



Formulas

Two loops
Junction law
Current in = current out
at(a) itotal = i1 +i2
Parallel arrangement
ΔV1= ΔV2
iTotal = ΔVBattery/Rtotal
iTotal = i1 +i2
ΔVBattery/Rtotal=ΔV1/R1+ΔV2/R2
1/Rtotal = 1/R1 + 1/R2
Rtotal = (1/R1 +1/R2)
R1=R2 =10Ohms
1/Rt = 1/10 + 1/10
= 2/10 = 1/5
RT= 5 ohms
Voltage loop law
One loop
ΔVbattery = Δ1+Δ2
Voltage drop of battery must equal
Sum of voltage drops around one loop.

Sponsored by **ApolloPad.com**
Everyone has a novel in them. Finish Yours!
https://apollopad.com