

Membrane Transport

Passive Transport: net movement of molecules from H to L concentration without ATP; used for import and export of materials

Facilitated Diffusion: (1) Large quantities of water pass through aquaporins (2) Charged ions, (Na⁺ and K⁺), require channel proteins to move through membrane (3) Membranes may become polarized by movement of ions across the membrane

Active Transport: uses ATP to transport molecules and establish/maintain concentration gradients; requires membrane proteins

Exocytosis: internal vesicles fuse with plasma membrane and secrete large macromolecules out of cell

Endocytosis: cell takes in macromolecules and particulate matter by forming new vesicles derived from plasma membrane

Selective Permeability

-Selective permeability is a direct consequence of membrane structure

-Small, non polar molecules can pass through (N₂, O₂ and CO₂)

-Hydrophilic substances (large polar molecules and ions) need embedded channels and transport proteins

-Polar uncharged molecules (H₂O) pass though in small amounts

-Allows for the formation of concentration gradients of solutes across the membrane

Surface Area to Volume Ratio

-Smaller cells typically have a higher SA:V for more efficient exchange of materials with environment

-As V increases, SA decreases, demand for internal resources increases


-Increasing cell size decreases SA:V

-Membrane folds can increase SA:V

Prokaryotic vs. Eukaryotic

Prokaryotes	Eukaryotes
Typically have circular chromosomes (plasmid)	Typically have multiple linear chromosomes (can have plasmids too)
Unicellular	Multicellular
No membrane bound nucleus	Membrane bound nucleus
Rare: microtubules, cytoskeleton; also chlorophyll scattered in cell	Lysosomes, peroxisomes, microtubules, ER, Mitochondria, Cytoskeleton, Vesicles, Golgi, Chloroplasts
Smaller ribosomes, have vacuoles	Larger ribosomes, have vacuoles
Chemically complex cell wall	Chemically simple cell wall
1-10um	10-100um
Groups of genes (operons) are transcribed in a single mRNA molecule	Groups of genes may be influenced by the same transcription factors to coordinately regulate expression

Cell Compartmentalization



-Membrane-bound organelles evolved from once free-living prokaryotic cells via endosymbiosis

-Prokaryotes generally lack internal membrane-bound organelles but have internal regions with specialized structures and functions.

-Eukaryotic cells maintain internal membranes that partition the cell into specialized regions

Cell Organelles

Ribosomes	Comprise ribosomal RNA (rRNA) and protein; Synthesize protein according to mRNA sequence; Found in all forms of life (evidence of common ancestor)
Endoplasmic Reticulum (ER)	Rough ER: compartmentalizes the cell; Smooth ER: detoxification and lipid synthesis
Golgi	Fold and chemically modify newly synthesized proteins; Packaging proteins for trafficking
Mitochondria	Powerhouse of cell; Double membrane provides compartments for different metabolic reactions



By PrincessB3ll3

Not published yet.

Last updated 14th May, 2020.

Page 1 of 2.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish Yours!


<https://apollopad.com>

Cell Organelles (cont)

Lysosomes Contain hydrolytic enzymes for intracellular digestion, recycling of a cell's organic materials, apoptosis

Vacuoles Storage and release of macromolecules and cellular waste products; In plants, aids in retention of water for turgor pressure

Osmosis



Water moves (by osmosis) from areas of H₂O potential/L solute concentration to areas of L H₂O potential/H solute concentration



By PrincessB3ll3

cheatography.com/princessb3ll3/

Not published yet.

Last updated 14th May, 2020.

Page 2 of 2.

Sponsored by [ApolloPad.com](https://apollopad.com)

Everyone has a novel in them. Finish

Yours!

<https://apollopad.com>