

### outlines

Oxidative phosphorylation (Electron transport chain + Chemiosmosis)

Eukaryotic Oxidative Phosphorylation Takes Place in Mitochondria

Oxidative Phosphorylation Depends on Electron Transfer (loss of electrons)

The Respiratory Chain Consists of Four Complexes: Three Proton Pumps and a Physical Link to the Citric Acid Cycle

A Proton Gradient Powers the Synthesis of ATP

Many Shuttles Allow Movement Across the Mitochondrial Membranes

The Regulation of Cellular Respiration Is Governed Primarily by the Need for ATP

Electron transfer potential of NADH and FADH<sub>2</sub> -> Phosphoryl transfer potential of ATP


### outline

A sedentary male of 70 kg (154 lbs) requires about 8400 kJ (2000 kcal) for a day's worth of activity.


To provide this much energy requires 83 kg of ATP. However, human beings possess only about 250 g of ATP at any given moment.

The disparity between the amount of ATP that we have and the amount that we require is compensated by recycling ADP back to ATP. Each ATP molecule is recycled approximately 300 times per day. This recycling takes place primarily through oxidative phosphorylation.


### bonding



### Overview of oxidative phosphorylation



### Eukaryotic Oxi\_Phos\_ take place in Mitochondria



### Mitochondria

Humans contain an estimated 14,000 m<sup>2</sup> of inner mitochondrial membrane.


The mitochondrial matrix is the site of most of the reactions of the citric acid cycle and fatty acid oxidation. In contrast, oxidative phosphorylation takes place in the inner mitochondrial membrane

The outer membrane is quite permeable to most small molecules and ions because it contains many copies of mitochondrial porin, a 30- to 35-kD pore-forming protein also known as VDAC, for voltage-dependent anion channel.

### Mitochondria (cont)

In contrast, the inner membrane is impermeable to nearly all ions and polar molecules.

### Electron transfer



### Electron transfer

Volt potential difference between NADH and O<sub>2</sub> drives electron transport and favors formation of a proton gradient

NADH->Pump1->CoQ->Pump3->Cyt c-->Pump4->2O<sub>2</sub>->2H<sub>2</sub>O

### Respiratory Chain have Four Complexes

Three Proton Pumps and a Physical Link to the Citric Acid Cycle

Electrons are transferred from NADH to O<sub>2</sub> through a chain of three large protein complexes called **NADH-Q oxidoreductase (I)**, **Q-cytochrome c oxidoreductase (III)**, and **cytochrome c oxidase (IV)**.

Electron flow within these transmembrane complexes leads to the transport of protons across the inner mitochondrial membrane.


A fourth large protein complex, called **succinate-Q reductase (II)**, contains the succinate dehydrogenase that generates FADH<sub>2</sub> in the citric acid cycle.

Oxidoreductase = reductase = dehydrogenase

Ubiquinone (Coenzyme Q) also carries electrons from FADH<sub>2</sub> (generated by citric acid cycle) generated through succinate-Q reductase



### Electrons flow down an energy gradient from NADH to O<sub>2</sub>



### Components of mitochondrial etc

Enzyme complex	Prosthetic group	Oxidant or reductant	
		Matrix side	Membrane core / Cytoplasmic side
NADH-Q oxidoreductase <b>Complexes I</b>	FMN Fe-S	NADH	Q
Succinate-Q reductase <b>Complexes II</b>	FAD Fe-S	Succinate	Q
Q-cytochrome c oxidoreductase <b>Complexes III</b>	Heme b <sub>L</sub> Heme b <sub>H</sub> Heme c <sub>1</sub> Fe-S		Cytochrome c
Cytochrome c oxidase <b>Complexes IV</b>	Heme a Heme a <sub>3</sub> Cu <sub>A</sub> and Cu <sub>B</sub>		Cytochrome c



By **rhettbro**  
[cheatography.com/rhettbro/](https://cheatography.com/rhettbro/)

Published 14th April, 2021.  
 Last updated 14th April, 2021.  
 Page 2 of 2.

Sponsored by **ApolloPad.com**  
 Everyone has a novel in them. Finish Yours!  
<https://apollopad.com>