

1 Sample T-Testing

For **numerical** data.

Compares a sample mean to a hypothetical population mean.

```
from scipy.stats import ttest_1samp
```

`ttest_1samp` requires two inputs, a distribution of values and an expected mean.

```
tstat, pval = ttest_1samp(example_distribution, expected_mean)
```

2 Sample T-Test

For **numerical** data.

Compares two sets of data, which are both approximately normally distributed.

The *null hypothesis*, in this case, is that the two distributions have the same mean.

```
from scipy.stats import ttest_ind
```

It takes the two distributions as inputs and returns the t-statistic and a p-value.

```
t, pval = ttest_ind(dataset1, dataset2)
```

ANOVA

For **numerical** data.

Compares more than two numerical datasets.

ANOVA (Analysis of Variance) tests the null hypothesis that all of the datasets have the same mean.

```
from scipy.stats import f_oneway
```

It takes in each dataset as a different input and returns the t-statistic and the p-value.

```
t, pval = f_oneway(a, b, c)
```

Tukey's Range Test

For **numerical** data.

We can perform a **Tukey's Range Test** to determine the difference between datasets.

```
from statsmodels.stats.multicomp import pairwise_tukeyhsd
```

We have to provide the function with one *list of all of the data* and a *list of labels* that tell the function which elements of the list are from which set. We also provide the *significance level* we want, which is usually 0.05.

```
values = np.concatenate([a, b, c])
```

```
labels = ['a'] * len(a) + ['b'] * len(b) + ['c'] * len(c)
```

```
tukey_results = pairwise_tukeyhsd(values, labels, 0.05)
```

Binomial Test

For **categorical** data.

To analyze a dataset with two different possibilities for entries.

The **null hypothesis**, in this case, would be that there is no difference between the observed behavior and the expected behavior.

```
from scipy.stats import binom_test
```

`binom_test` requires three inputs, the number of observed successes, the number of total trials, and an expected probability of success.

```
pval = binom_test(525, n=1000, p=0.5)
```

Chi Square Test

For **categorical** data.

To compare two or more categorical datasets.

```
from scipy.stats import chi2_contingency
```

The input to `chi2_contingency` is a **contingency table** where:

- **The columns** represent different outcomes, like "Survey Response A" vs. "Survey Response B" or "Clicked a Link" vs. "Didn't Click"

- **The rows** are each a different condition, such as men vs. women or Interface A vs. Interface B

```
X = [[30, 10],
```

```
     [35, 5],
```

```
     [28, 12],
```

```
     [20, 20]]
```

```
_, pval, _, _ = chi2_contingency(X)
```



By **sasha2411** (sasha2411)

Published 26th July, 2018.

Last updated 26th July, 2018.

Page 1 of 1.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>