

Trig Integrals

$\int \sin x \, dx$	$= -\cos x + C$
$\int \cos x \, dx$	$= \sin x + C$
$\int \sec^2 x \, dx$	$= \tan x + C$
$\int \tan x \, dx$	$= \ln \sec x + C$
$\int \sec x \tan x \, dx$	$= \sec x + C$
$\int \csc^2 x \, dx$	$= -\cot x + C$
$\int \csc x \cot x \, dx$	$= -\csc x + C$
$\int \cot x \, dx$	$= \ln \sin x + C$

Trig Identities

$$\int (1/x^2 + a^2) \, dx = 1/a \tan^{-1}(x/a) + C$$

$$\int (1/\sqrt{a^2 - x^2}) \, dx = \sin^{-1}(x/a) + C$$

(While $a > 0$)

Area Between Curves

$$\text{Area} = \int [\text{Height}] \text{Width}$$

$$A = \int (f(x) - g(x)) \, dx$$

1. Graph Equations
2. Label
3. Determine how to slice
4. Set up dA
5. $dA = \text{height} \cdot dx$
6. Get range a & b from intersections
7. Plug in and find area

There is never (-) area.

Volume by Disk

$$dV = A(x) \, dx$$

$$V = \int A(x) \, dx$$

$$\text{Volume} = \int \text{Radius}^2 \cdot \text{Thickness}$$

$$V = \int (\pi(r)^2) \, dx$$

Volume by Washer

$$dV = A(x) \, dx$$

$$V = \int A(x) \, dx$$

$$\text{Volume} = \int [(\pi r_{\text{out}}^2) - (\pi r_{\text{in}}^2)] \, dx$$

Slice Perpendicular to Axis of Rotation

Volume by Shell

$$d\text{Volume} = \text{Circumference} \cdot d\text{Area}$$

$$dV = (2 \pi r) \, d\text{Area}$$

$$V = \int (2 \pi r) (\text{Area}) \, dx$$

1. Write: $dV = 2 \pi r \, dA$
2. Find $dA(\text{height} \, dx)$
3. Find Radius (x or y)
4. Plug in
5. Take integral

Slice Parallel to Axis of Rotation

Average Value of a Function

$$\text{Average Value} = 1/(b-a) \cdot \int f(x) \, dx$$

Symmetry:

If $f(x)$ is EVEN, then $\int f(x) \, dx$ from $-a$ to $a = 2 \int f(x)$ from 0 to a

If $f(x)$ is ODD, then $\int f(x) \, dx$ from $-a$ to $a = 0$

Important Integrals

$$\int c \, f(x) \, dx = c \int f(x) \, dx$$

$$\int [f(x) + g(x)] \, dx = \int f(x) \, dx + \int g(x) \, dx$$

$$\int 1/x \, dx = \ln|x| + C$$

$$\int e^x \, dx = e^x + C$$

$$\int b^x \, dx = (b^x / \ln b) + C$$

Methods of Integration

Method	When to Use	Example
U-Substitution	When a Polynomial is raised to a power > 1	$\int (3x + 5)^5$
Integration by Parts	When U-Sub will not work	$\int x e^x$
Trigonometric Integration	Only Trig raised to powers	$\int \sin^6 x \cos^3 x \, dx$
Trigonometric Substitution	$3/2$ powers or $\sqrt{a^2 - x^2}$ etc.	$dx / (x^2 \sqrt{25 - x^2})$

U-Substitution



By smithandrewa

Not published yet.

Last updated 27th February, 2019.

Page 1 of 2.

Sponsored by [Readable.com](https://readable.com)

Measure your website readability!

<https://readable.com>

Integration by Parts

Logarithmic

Inverse trig

Algebraic

Trigonometric

Exponential

$$\int u \, dv = u \, v - \int v \, du$$

1. Write $u \, v - \int v \, du$

2. Use LIATE to find u ; the other term becomes dv

3. Setup $u = dv = du = v =$

4. Solve

Cyclical Functions will need to be split and substituted.

Trigonometric Integration

Identities

$$\sin^2 t + \cos^2 t = 1$$

$$\sin^2 t = \frac{1}{2} [1 - \cos(2t)]$$

$$\cos^2 t = \frac{1}{2} [1 + \cos(2t)]$$

Can use with U-Substitution

Don't change all of the trig to the same form.

Trigonometric Integration

Identities

$$\sin^2 t + \cos^2 t = 1$$

$$\sin^2 t = \frac{1}{2} [1 - \cos(2t)]$$

$$\cos^2 t = \frac{1}{2} [1 + \cos(2t)]$$

Can use with U-Substitution

Don't change all of the trig to the same form.

Trigonometric Substitution

Pythag. Identities

$$\sin^2 + \cos^2 = 1$$

$$1 + \tan^2 = \sec^2$$

$$1 + \cot^2 = \csc^2$$

1. Identify a and u

2. Sub in the trig

3. Manipulate to simplify

4. Get rid of trig with a triangle



By **smithandrewa**

cheatography.com/smithandrewa/

Not published yet.

Last updated 27th February, 2019.

Page 2 of 2.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>