

### ACUTE DISEASES OF THE NEWBORN

- high-risk neonate regardless of Gestational Age
- begins 23wks - 28 days post birth

### CLASSIFICATIONS OF HIGH RISK NB

#### According to Size

- Low Birth Weight **LBW** (<2.5kg)
- Very Low Birth Weight **VLBW** (<1.5kg)
- Extremely Low Birth Weight **ELBW** (<1kg)
- Appropriate for Gestational Age **AGA** (10%-90%)
- Small for Gestational Age **SGA** (<10%)
- Large for Gestational Age **LGA** (>90%)
- Intrauterine Growth Restriction **IUGR**

#### >Risk Factors:

- Hereditary
- Placental Insufficiency
- Maternal Disease


#### According to Gestational Age (regardless of BW)

- Late Preterm (**34-36wks AOG**)
- Preterm (**<37wks AOG**)
- Full term (**38-42wks AOG**)
- Post term (**>42wks AOG**)

#### According to Mortality

- Live Birth
- Fetal Death
  - = death before 20wks
- Neonatal Death
  - = death within first 27wks of extra-uterine life
- Perinatal Mortality
  - = total # of fetal & neonatal death/1000 live births
- Postnatal Death
  - = death 28 days - 1y/o

### Intrauterine Growth Curve



### ASSESSMENT OF HIGH-RISK NB

#### 1. Physical Assessment

- General Assessment
  - > BW
  - > Anthropometric Measurements
  - > Deformities
  - > Signs of distress (poor color, mottling, hypotonia)

#### 2. Respiratory Assessment

- Chest Shape (barrel/concave)
- Describe use of accessory muscles
- Determine RR; O<sub>2</sub> Sat
- Auscultation

#### 3. Cardiovascular Assessment

- HR and rhythm
- Auscultation
- Determine Point of Maximal Impulse (PMI)
- Color
  - > mucous membranes, lips, BP, perfusion

#### 4. Genitourinary Assessment

- Genitalia and abnormalities
- Describe urine

### ASSESSMENT OF HIGH-RISK NB (cont)

> amount, pH, specific gravity

#### 5. Gastrointestinal Assessment

- Presence of abdominal distention, regurgitation
- Stool assessment
  - > amount, color, consistency

#### 6. Neurologic-Musculoskeletal Assessment

- Movements, Level of Activity with stimulation
- Changes in Head Circumference

#### 7. Temperature

- Determine axillary temperature

### HIGH-RISK CONDITIONS RT DYSPMAT-URITY

#### 1. Preterm Infants

- Cause:
  - > idiopathic
- Risk Factors:
  - > low socio-economic status
  - > multigravida
  - > gestational HTN
- Characteristics:
  - > very small and thin; little SQ fat
  - > proportionally large head
  - > bright pink, shiny, smooth skin
  - > abundant fine lanugo
  - > ear cartilage soft and pliable
  - > male NB = few scrotal rugae; cryptochordism
  - > female NB = labia minora & clitoris prominent

#### 2. Post-term Infants

- Cause:
  - > idiopathic
- Characteristics:
  - > absent lanugo

### HIGH-RISK CONDITIONS RT DYSMAT-URITY (cont)

- > abundant scalp hair; long fingernails
- > cracked skin/parchment-like/desquamation
- > depleted SQ fat
- > little vernix caseosa

### PROBLEMS RT GESTATIONAL WEIGHT

### SGA RT Intrauterine Growth Restriction (IUGR)

- **Cause:**
  - > poor nutrition
  - > adolescent pregnancy
  - > placental anomaly
  - > maternal systemic disease (HTN, DM)
- **Diagnostic Evaluation**
  - > fundal height < expected
  - > UTZ = ↓ size; placental grading; amniotic fluid
    - > biophysical profile
    - > non-stress test (NST)
- **Fetal Implications**
  - > poor skin turgor
  - > large head, small body
  - > small liver
  - > skull sutures widely separated
  - > ↑ Hct level
  - > polycythemia (↑ RBC)
  - > hypoglycemia (<45mg/dL)

### LGA (Macrosomia)

- appears healthy, but will soon reveal underdevelopment
- **Causes:**
  - > gestational DM (GDM)
  - > multiparity
  - > **Beckwith Syndrome** (overgrowth+macroglossia)
    - > congenital anomalies (omphalocele)
- **Diagnostic Evaluation**
  - > UTZ
  - > NST
  - > amniocentesis
- **Fetal Implications**
  - > immature reflexes
  - > extensive bruising/birth injury **Erb-Duchenne**
    - > caput succedaneum; cephalhematoma
    - > hyperbilirubinemia
    - > polycythemia vera
    - > cyanosis
    - > ↑ insulin (up to 24hrs post birth=hypoglycemia)

### MANAGEMENT OF HIGH-RISK NEWBORN

#### NEWBORN PRIORITIES

1. **Initiating/Maintaining Respiration**
  - most deaths occur within 48hrs
  - ineffective respiration = cerebral hypoxia
    - > Management:
      - O2 administration
      - appropriate positioning to open airway
        - resuscitation+ventilation
2. **Establish Extraterine Circulation**
  - > Management:

### MANAGEMENT OF HIGH-RISK NEWBORN (cont)

- closed-chest massage (1-2cm, 100x/min)
  - lung ventilation (30x/min)
  - monitor pulse oximeter
  - 0.1-0.3mL/kg **Ephinephrine** may be sprayed on ET tube
    - transfer to NICU
- 3. **Maintain Fluid Balance**
  - > Management: (after initial resuscitation)
    - Hypoglycemia (**D10W IVF**)
    - Hypotension (vasopressor)
- Dopamine**
  - Hypovolemia (**NSS/RL IVF**)
  - Dehydration (**RL/D5W IVF**)
- 4. **Maintaining Thermoneutrality**
  - > Management:
    - thorough drying
    - skin-skin contact
    - neutral thermal environment
- 5. **Establishing Adequate Nutritional Intake**
  - > Management:
    - parenteral/enteral nutrition
    - breastfeeding
- 6. **Establishing Waste Elimination**
  - Immature infants void within 24hrs
  - stool passage may be later than term infants
- 7. **Protection from Infection**
  - > Prevention:
    - handwashing and PPE use
    - standard precautions
    - physical isolation
- 8. **Skin Care**
  - ↑ skin sensitivity & fragility



By vcroosevelt (vcroosevelt)

[cheatography.com/vcroosevelt/](https://cheatography.com/vcroosevelt/)

Not published yet.

Last updated 5th April, 2023.

Page 2 of 4.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>

### MANAGEMENT OF HIGH-RISK NEWBORN (cont)

- > Management:
  - Zinc Oxide-based tape is used
  - avoid use of solvents

#### 9. Establishing Mother-Infant Bonding

- parents kept informed
- spend time with NB

\*2. (1:3 = Lung ventilation:Cardiac massage)

\*3. Monitor UO (if UO=<2mL/kg/hr = inadequate fluid intake)

\*4. 3 Main Methods for Neutral Thermal Environment: Incubator, Radiant panel, Bassinet

\*5. If gavage fed, provide oral stimulation to develop effective sucking reflex

### ACUTE CONDITIONS OF NEONATES

#### Respiratory Distress Syndrome (RDS)

- Hyaline Membrane Disease
- surfactant deficiency
- **Types:**
  - > *Structural*
    - lungs are underdeveloped
    - respiratory muscle prone to fatigue
  - > *Functional*
    - deficient surfactant
- **Risk Factors:**
  - > Multifetal pregnancy
  - > GDM
  - > CS Delivery
  - > Cold stress
  - > Asphyxia
  - > Hx of RDS
- RDS of Non-Pulmonary Origin Risk Factors:
  - > Sepsis
  - > Cardiac Defect
  - > Hypoglycemia
  - > Metabolic Acidosis

#### Respiratory Distress Syndrome (RDS) (cont)

- > Drugs
- **Clinical Manifestations:**
  - > tachypnea (>60cpm)
  - > retractions; nasal flaring
  - > inspiratory crackles
  - > circumoral and central cyanosis
- **Laboratory Diagnoses:**
  - > Glucometry (tests hypoglycemia)
  - > ABG (tests acidosis, hypoxia, hypercapnia)
  - > CXR
    - diffuse granular pattern = alveolar atelectasis
    - dark streaks = dilated, air-filled bronchioles
- **Treatment:**
  - > ventilation and oxygenation with Continuous Positive Airway Pressure (CPAP)
  - > maintain acid-base balance
  - > neutral thermal environment
  - > maintain hydration and electrolytes
  - > avoid nipple and gavage feedings
  - > administer exogenous surfactants
- **Nursing Responsibilities:**
  - > collect and monitor ABG
  - > O2 monitoring
  - > assess tolerance on procedure/drug

\* Surfactants produced at 24wks AOG, matures at 36wks

\* Surfactant Complications: pulmonary hemorrhage; mucus plugging

#### Meconium Aspiration Syndrome

##### Meconium

- sticky and tarlike; present at bowel 10wks AOG

- accumulates at 16wks AOG

##### Meconium Aspiration

- occurs inside utero/at first breath at birth

- occurs when the vagus reflex is stimulated due to hypoxia → releasing meconium to amniotic fluid

- NB born at breech position

##### - Pathophysiology:

> hypoxia → meconium passing → aspiration → obstruction → atelectasis → respiratory failure

##### - Clinical Manifestations:

- > tachypnea; retractions; expiratory grunting; nasal flaring
- > cyanosis/pallor
- > barrel chest (from hyperinflation)
- > hypoglycemia; hypocalcemia

##### - Diagnostic Evaluation:

- > laryngoscopy
- > CXR

##### - Management:

- > tracheal suctioning
- > intubation (in severe cases)
- > surfactant administration
- > Echocardiography (diagnose shunting)
- > chest physiotherapy

### Apnea of Prematurity (AOP)

#### Apnea

- cessation of respiration that lasts >20secs, accompanied by bradypnea and cyanosis

#### Types:

##### 1. Central Apnea

- absent function of diaphragmatic and other respiratory muscles  
- CNS does not transmit signals to respiratory muscles

##### 2. Obstructive Apnea

- airflow stops due to obstruction

##### 3. Mixed Apnea

- central + obstructive  
- most common on premature infants

#### - Causes:

- > prematurity (weak thorax muscles)
- > airway obstruction
- > anemia; polycythemia vera
- > hypoglycemia; hypocalcemia
- > sepsis; meningitis; seizures

#### - Management:

- > gentle tactile stimulation
- > Caffeine Citrate PO/Parenteral (CNS Stimulant)
  - > monitor weight and UO (Caffeine citrate = diuretic)
  - > nasal CPAP & nasal intermittent positive pressure ventilation
  - > neutral thermal environment

#### - Nursing Responsibilities:

- > routine observation (RR & HR)

### Apnea of Prematurity (AOP) (cont)

- > gentle tactile stimulation, if it fails, raise chin to open airway
- > careful burping = reduces apnea
- > never take rectal temperature



By **vcroosevelt** (vcroosevelt)

[cheatography.com/vcroosevelt/](https://cheatography.com/vcroosevelt/)

Not published yet.

Last updated 5th April, 2023.

Page 4 of 4.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>